

11/05/99
JCS66 U.S. PTO

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. B0801/7159(ERP)

First Named Inventor or Application Identifier

BAND, Hamid et al.

Express Mail Label No. EL056790199US

Date of Deposit November 5, 1999

11/05/99
JCS66 U.S. PTO
19/434708

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents

ADDRESS

Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

TO:

1. Fee Transmittal Form
(Submit an original, and a duplicate for fee processing)
2. X Specification [Total pages 64]
46 pages specification; 9 pages Seq. Listing
1 pages abstract
8 pages claims
69 claims
3. X Drawing(s) (35 USC 113) [Total sheets 9]
X Informal ☐ Formal [Total drawings 8, including Figs. 1A, 1B, 2, 3A, 3B, 3C, 3D, 4A, 4B, 5A, 5B, 6, 7, and 8]
4. ☐ Oath or Declaration [Total pages]
 - a. ☐ Newly executed (original or copy)
 - b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
[Note Box 5 below]
 - i. ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) and 1.33(b).
5. ☐ Incorporation by Reference
(usable if Box 4b is checked)
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Microfiche Computer Program (Appendix)
7. X Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - a. X Computer Readable Copy
 - b. X Paper Copy (identical to computer copy)
 - c. X Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

8. ☐ Assignment Papers (cover sheet & documents(s))
9. ☐ 37 CFR 3.73(b) Statement ☐ Power of Attorney
(when there is an assignee)
10. ☐ English Translation of Document (if applicable)
11. ☐ Information Disclosure ☐ Copies of IDS
Statement (IDS)/PTO-1449 Citations
12. X Preliminary Amendment
13. X Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
14. ☐ Small Entity ☐ Statement filed in prior
Statement(s) application, Status still proper
and desired
15. ☐ Certified Copy of Priority Document(s)
(if foreign priority is claimed)
16. Other: This application claims priority under Title 35 §119(e), of United States Provisional Application No. 60/107,470, filed November 6, 1998, and entitled CHARACTERIZATION OF NOVEL GENE CBL-SL, the entire contents of which are incorporated herein by reference.

17. If a CONTINUING APPLICATION, check appropriate box and supply the requisite information:


- ☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.:
- ☐ Cancel in this application original claims of the prior application before calculating the filing fee.
- ☐ Amend the specification by inserting before the first line the sentence:

This application is a ☐ continuation ☐ divisional of application serial no. , filed , entitled , and now .

18. CORRESPONDENCE ADDRESS*Correspondence address below*

ATTORNEY'S NAME	Elizabeth R. Plumer, Reg. No. 36,637				
NAME	Wolf, Greenfield & Sacks, P.C.				
ADDRESS	600 Atlantic Avenue				
CITY	Boston	STATE	MA	ZIP	02210
COUNTRY	USA	TELEPHONE	(617) 720-3500	FAX	(617) 720-2441

19. SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT REQUIRED

NAME	Elizabeth R. Plumer, Reg. No. 36,637
SIGNATURE	
DATE	November 5, 1999

0944703-1099
BOSTON 80242460


ATTORNEY'S DOCKET NO.: B0801/7159(ERP)

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: Hamid BAND et al.
Serial No: Not Yet Assigned
Filed: Herewith
For: CHARACTERIZATION OF NOVEL GENE CBL-SL
Examiner: Not Yet Assigned
Art Unit: Not Yet Assigned

CERTIFICATE OF MAILING UNDER 37 C.F.R. §1.8(a)

The undersigned hereby certifies that this document is being placed in the United States mail with first-class postage attached, addressed to Box: Patent Application, Assistant Commissioner for Patents, Washington, D.C. 20231, on the 5th day of November, 1999.


Elizabeth R. Plumer

Box: Patent Application
Assistant Commissioner for Patents
Washington, D.C. 20231

PRELIMINARY AMENDMENT


Sir:

Before calculating the fees, please amend the above-identified application as follows:

Please cancel claims 8, 10, 13-16, 18-20, 23-24, 26-33, 35-49, 51, 55-59, and 62-69.

Respectfully submitted,

WOLF, GREENFIELD & SACKS, P.C.


Elizabeth R. Plumer, Reg. No.: 36,637
WOLF, GREENFIELD & SACKS, P.C.
600 Atlantic Avenue
Boston, MA 02210-2211
Attorneys for Applicant(s)
Tel. No. (617) 720-3500

ATTORNEY'S DOCKET NO.: B0801/7159

Date: November 5, 1999

x11/6/99

CHARACTERIZATION OF NOVEL GENE CBL-SL

Related Applications

5 This application claims priority under 35 U.S.C. §119 from Provisional U.S. Patent Application Serial No. 60/107,470 filed on November 6, 1998, entitled CHARACTERIZATION OF NOVEL GENE CBL-SL. The contents of the provisional application are hereby expressly incorporated by reference.

Government Support

10 This work was funded in part by grant numbers RO1-CA75075-02 and RO1-CA76118-01 from the National Institutes of Health, and IM-79014 from the American Cancer Society. Accordingly, the United States Government may have certain rights to this invention.

Field of the Invention

15 This invention relates to nucleic acids and encoded polypeptides of cbl-SL, and diagnostics and therapeutics related to medical conditions associated with such genes and polypeptides, including cancers.

Background of the Invention

20 Tyrosine phosphorylation provides a universal mechanism of signal transduction in response to extracellular cues that regulate proliferation and differentiation in normal cells. Uncontrolled tyrosine kinase activation is implicated in proliferation of cancerous cells and deficiencies of specific tyrosine kinases result in a number of pathological conditions such as developmental abnormalities or immuno-deficiencies. Little is known at present about how cellular tyrosine kinases are regulated. Genetic studies in *C. elegans* have identified the *sli-1* proto-oncogene product as a potential regulator of tyrosine kinase signaling.

25 *Sli-1* is a member of the Cbl family of proto-oncogenes that includes c-Cbl, Cbl-b and the *Drosophila* gene D-Cbl. Recent studies report that Cbl, the 120 kDa protein product of the c-cbl proto-oncogene (first identified as part of a transforming gene of a murine retrovirus and whose expression is predominant in haematopoietic cells), consists of an amino-terminal transforming region, a zinc ring finger, multiple proline-rich stretches, and several potential phosphotyrosine-containing motifs.

30 Cbl is reportedly tyrosine-phosphorylated, at a fast rate, in response to stimulation of a variety of cell-surface receptors and upon integrin-mediated cell adhesion. Cbl becomes associated with a number of intracellular signalling molecules such as protein tyrosine kinases, phosphatidylinositol-3-kinase, Crk, 14-3-3 proteins, and other cytoskeletal and adaptor proteins through different protein-interacting modules (including Src Homology 2 and

3 -i.e., SH2 and SH3- domains), leading to the formation of multimolecular signalling complexes. Cbl can be rendered into a transforming factor by engineering mutations into its amino acid sequence. Cbl and its transforming mutants have been reported to display both negative and positive regulatory activities in protein tyrosine kinase- and Ras-mediated signalling pathways. The N-terminal region of c-Cbl (c-Cbl-N) harbors a tyrosine kinase binding domain, which presumably plays a regulating role in c-Cbl's transforming capabilities.

There exists a need to identify agents that desirably influence a cell's growth, differentiating and proliferative characteristics, and to provide diagnostics, research tools and therapeutics relating to such novel agents.

These and other objects will be described in greater detail below.

Summary of the Invention

We describe herein the molecular cloning and characterization of cbl-SL, a novel molecule that is believed to play a role in the regulation of a cell's growth, differentiation and proliferation. Abberant expression levels and mutant forms of tyrosine kinases which are the likely targets of cbl-SL have been found in patients with certain types of cancer, suggesting that cbl-SL may play a role in cell cycle control and cancerous cell proliferation in particular.

The invention provides isolated nucleic acid molecules, unique fragments of those molecules, expression vectors containing the foregoing, and host cells transfected with those molecules. The invention also provides isolated binding polypeptides and binding agents which bind such polypeptides, including antibodies. The foregoing can be used, *inter alia*, in the diagnosis or treatment of conditions characterized by the aberrant expression levels and/or the presence of mutant forms of a cbl-SL nucleic acid or polypeptide. The invention also provides methods for identifying pharmacological agents useful in the diagnosis or treatment of such conditions.

According to one aspect of the invention, isolated nucleic acid molecules that code for a cbl-SL polypeptide are provided and include: (a) nucleic acid molecules which hybridize under stringent conditions to a molecule consisting of a nucleic acid of SEQ ID NO:1 and which code for a cbl-SL polypeptide, (b) deletions, additions and substitutions of (a) which code for a respective cbl-SL polypeptide, (c) nucleic acid molecules that differ from the nucleic acid molecules of (a) or (b) in codon sequence due to the degeneracy of the genetic code, or (d) complements of (a), (b) or (c). In certain embodiments, the isolated nucleic acid molecule comprises nucleotides 1-1547 of SEQ ID NO:1. In some embodiments the isolated

nucleic acid molecules are those comprising the human cDNA or gene corresponding to SEQ ID NO:3. The isolated nucleic acid molecule also can comprise a molecule which encodes the polypeptide of SEQ ID NO:2 and has cbl-SL binding activity.

The invention in another aspect provides an isolated nucleic acid molecule selected from the group consisting of (a) a unique fragment of nucleic acid molecule of SEQ ID NO:1 (of sufficient length to represent a sequence unique within the human genome), (b) complements of (a), provided that the fragment includes a sequence of contiguous nucleotides which is not identical to a sequence selected from the sequence group consisting of (1) sequences having the GenBank and EMBL accession numbers of Table I, (2) complements of (1), and (3) fragments of (1) and (2).

In one embodiment, the sequence of contiguous nucleotides is selected from the group consisting of (1) at least two contiguous nucleotides nonidentical to the sequence group, (2) at least three contiguous nucleotides nonidentical to the sequence group, (3) at least four contiguous nucleotides nonidentical to the sequence group, (4) at least five contiguous nucleotides nonidentical to the sequence group, (5) at least six contiguous nucleotides nonidentical to the sequence group, (6) at least seven contiguous nucleotides nonidentical to the sequence group.

In another embodiment, the fragment has a size selected from the group consisting of at least: 8 nucleotides, 10 nucleotides, 12 nucleotides, 14 nucleotides, 16 nucleotides, 18 nucleotides, 20 nucleotides, 22 nucleotides, 24 nucleotides, 26 nucleotides, 28 nucleotides, 30 nucleotides, 40 nucleotides, 50 nucleotides, 75 nucleotides, 100 nucleotides, 200 nucleotides, 1000 nucleotides and every integer length therebetween.

According to another aspect, the invention provides expression vectors, and host cells transformed or transfected with such expression vectors, comprising the nucleic acid molecules described above.

According to another aspect of the invention, an isolated polypeptide is provided. The isolated polypeptide is encoded by the foregoing isolated nucleic acid molecules of the invention. In some embodiments, the isolated polypeptide is encoded by the nucleic acid of SEQ ID NO:1, giving rise to a ~50 kd polypeptide having the sequence of SEQ ID NO:2 that can associate with a tyrosine kinase to form a complex, and thereby could regulate a cell's growth, differentiation and proliferation. In other embodiments, the isolated polypeptide may be a fragment or variant of the foregoing of sufficient length to represent a sequence unique within the human genome, and identifying with a polypeptide that can associate with a tyrosine kinase to form a complex, and regulate a cell's growth, differentiation and

proliferation, provided that the fragment includes a sequence of contiguous amino acids which is not identical to any sequence encoded for by the nucleic acid sequence selected from the group consisting of SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9. In another embodiment, immunogenic fragments of the polypeptide molecules described above are provided.

According to another aspect of the invention, isolated binding polypeptides are provided which selectively bind a polypeptide encoded by the foregoing isolated nucleic acid molecules of the invention. Preferably the isolated binding polypeptides selectively bind a polypeptide which comprises the sequence of SEQ ID NO:2, SEQ ID NO:10, or fragments thereof, and that do not recognize ("cross-react") with epitopes from polypeptides encoded for by the nucleic acid sequence selected from the group consisting of SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9. In preferred embodiments, the isolated binding polypeptides include antibodies and fragments of antibodies (e.g., Fab, F(ab)₂, Fd and antibody fragments which include a CDR3 region which binds selectively to the cbl-SL polypeptide). In certain embodiments, the antibodies are human.

The invention also contemplates kits comprising a package including assays for cbl-SL epitopes, cbl-SL nucleic acids, and instructions, and optionally related materials such as controls, for example, a number, color chart, or an epitope of the expression product of the foregoing isolated nucleic acid molecules of the invention, for comparing the level of cbl-SL polypeptides or cbl-SL nucleic acid forms (wild-type or mutant) in a test sample to the level in a control sample. This comparison can be used to assess in a subject a risk of developing a cancer. The kits may also include assays for other known genes, and expression products thereof, associated with cancers (e.g., BRCA, p53, etc.).

According to another aspect of the invention, a method of screening for the presence of a carcinoma in a subject suspected of having a carcinoma is provided. The method involves (a) characterizing cbl-SL nucleic acid sequences in a test sample, wherein the test sample is obtained from a tissue of the subject, and (b) comparing the cbl-SL nucleic acid sequences of the test sample to cbl-SL nucleic acid sequences of a control sample (wild-type or mutant). An observed alteration or match in a cbl-SL nucleic acid sequence in the test sample as compared to cbl-SL nucleic acid sequences in the control sample, is indicative of the presence of carcinoma in the subject. In certain embodiments, the observed alteration is apparent when a cbl-SL nucleic acid sequence in the test sample is compared to wild-type cbl-SL nucleic acid sequences in the control sample. In other embodiments, the observed match is apparent

when a cbl-SL nucleic acid sequence in the test sample is compared to mutant cbl-SL nucleic acid sequences in the control sample.

In yet other embodiments, cbl-SL mRNA molecules are compared. Alteration of cbl-SL mRNA may be detected by hybridization of test sample mRNA to any of the foregoing cbl-SL nucleic acids of the invention.

In other embodiments, cbl-SL cDNA sequences are compared. The comparison is performed by hybridization of a cbl-SL cDNA probe to genomic DNA isolated from the test sample. In further embodiments, the genomic DNA is isolated from a non-neoplastic tissue of the subject and is subjected to Southern hybridization with a cbl-SL cDNA probe. The hybridizations are compared to (i) the cbl-SL cDNA probe to the test sample and/or (ii) the cbl-SL cDNA probe to the non-neoplastic tissues. The cbl-SL cDNA probe may detect a restriction fragment length polymorphism.

In yet other embodiments, cbl-SL nucleic acid sequences are compared, and the comparing is performed by determining the sequence of all or part of a cbl-SL cDNA in the test sample using a polymerase chain reaction. Deviations in the cbl-SL cDNA determined from that of the wild-type cbl-SL nucleic acid sequence shown in SEQ ID NO: 1, is indicative of the presence of the carcinoma in the subject.

In certain embodiments, the alteration of cbl-SL nucleic acid sequences is detected by identifying a mismatch between molecules (a) a cbl-SL cDNA or cbl-SL mRNA isolated from the tissue and (b) a nucleic acid probe complementary to the human wild-type cbl-SL nucleic acid sequence, when molecules (a) and (b) are hybridized to each other to form a duplex.

In certain other embodiments, cbl-SL nucleic acid sequences are compared and the alteration of cbl-SL nucleic acid sequences is detected by (a) amplifying cbl-SL cDNA sequences in the test sample, and (b) hybridizing the amplified cbl-SL cDNA sequences to nucleic acid probes which comprise cbl-SL sequences.

In yet other embodiments, cbl-SL nucleic acid sequences are compared, and the alteration of cbl-SL nucleic acid sequences is detected by molecular cloning of cbl-SL sequences in the test sample and characterized by sequencing all or part of the cloned cbl-SL gene.

In any of the foregoing embodiments, detection of alteration of cbl-SL nucleic acid sequences includes screening for a deletion mutation, a point mutation, and/or an insertion mutation. In preferred embodiments, the test sample may be obtained from tissues of a subject that include blood, breast, colon and prostate.

09434709 1059 66501 8024460

Another aspect of the invention is a method for determining the level of cbl-SL expression in a subject. Expression is defined either as cbl-SL mRNA expression or cbl-SL polypeptide expression. Various methods can be used to measure expression. Preferred embodiments of the invention include PCR and Northern blotting for measuring mRNA expression, and monoclonal or polyclonal cbl-SL antisera as reagents to measure cbl-SL polypeptide expression. In certain embodiments, test samples such as biopsy samples, and biological fluids such as blood, are used as test samples. Cbl-SL expression in a test sample of a subject is compared to cbl-SL expression in control sample to, e.g., assess the presence or absence or stage of a cancer in a subject.

The invention in another aspect involves a method for increasing cbl-SL expression in a subject that expresses a mutant cbl-SL. An isolated cbl-SL nucleic acid molecule of the invention or an expression product thereof is administered to a subject expressing a mutant cbl-SL, in an amount effective to increase wild-type cbl-SL expression in the subject.

Another aspect of the invention provides compositions comprising any of the foregoing isolated nucleic acid molecules of the invention, or expression products thereof, and which increase expression of cbl-SL (wild-type), and a pharmaceutically acceptable carrier.

According to another aspect, a method for downregulating (or inhibiting) expression of a tyrosine kinase in a cell is provided. The method involves contacting a cell expressing a tyrosine kinase with a cbl-SL polypeptide, in an amount effective to downregulate expression of the tyrosine kinase in the cell. Preferred cbl-SL polypeptides include the polypeptides of claim 12. In certain embodiments, the tyrosine kinase is selected from the group consisting of a receptor tyrosine kinase, a non-receptor tyrosine kinase, and any of the foregoing tyrosine kinases in complex with another agent. In one embodiment, the tyrosine kinase is ligand activated. In any of the foregoing embodiments, the tyrosine kinase can be phosphorylated. In certain embodiments, the cbl-SL and tyrosine kinase further complexes with an intermediate adapter molecule (e.g., Grb-2).

According to still another aspect of the invention, a method is provided for identifying lead compounds for an agent useful in the diagnosis or treatment of disease associated with cbl-SL activity. The method involves forming a mixture of a cbl-SL polypeptide, a tyrosine kinase that binds a cbl-SL polypeptide (such as the EGF receptor in the presence or absence of other molecules, e.g. Grb-2), and a candidate agent. The mixture is incubated under conditions which permit specific binding of the tyrosine kinase that binds a cbl-SL polypeptide to a cbl-SL polypeptide (i.e., in the absence of a candidate agent). A reference specific binding of the tyrosine kinase that binds a cbl-SL polypeptide to a cbl-SL polypeptide

is then detected, the reference specific binding being indicative of cbl-SL activity. The reference specific binding is then compared to a control. Detection of an increase in the foregoing activity relative to the reference specific binding in the presence of the candidate agent indicates that the candidate agent is a lead compound for an agent which increases the cbl-SL activity. Detection of a decrease in the foregoing activities relative to the reference specific binding in the presence of the candidate agent indicates that the candidate agent is a lead compound for an agent which decreases cbl-SL activity. In some embodiments, the binding can occur intracellularly and in a single cell. Preferred cbl-SL polypeptides include the polypeptides of claim 12. In certain embodiments, the tyrosine kinase is selected from the group consisting of a receptor tyrosine kinase, a non-receptor tyrosine kinase, and any of the foregoing tyrosine kinases in complex with another agent. In one embodiment, the tyrosine kinase is ligand activated. In any of the foregoing embodiments, the tyrosine kinase can be phosphorylated. In certain embodiments, the cbl-SL and tyrosine kinase further complexes with an intermediate adapter molecule (e.g., Grb-2).

The present invention thus involves, in several aspects, cbl-SL polypeptides, nucleic acids encoding those polypeptides, functional modifications and variants of the foregoing, useful fragments of the foregoing, as well as therapeutics and diagnostics relating thereto.

These and other aspects of the invention, as well as various advantages and utilities, will be more apparent with reference to the detailed description of the preferred embodiments.

Brief Description of the Drawings

Figure 1 shows side-by-side comparison of cbl (Fig. 1A) and cbl-SL (Fig. 1B) mRNA expression on identical Tissue Blots.

Figure 2 shows a cbl-SL mRNA expression in a Northern Blot.

Figure 3A and 3B show the specific binding of cbl-SL polyclonal antisera using Western blots. In Fig. 3C, an immunizing peptide was used as a competitor. Fig. 3D also shows the specific binding of cbl-SL polyclonal antisera on HA-purified protein lysates.

Figure 4A depicts a ~50kd cbl-SL protein band in a Western blot. Fig. 4B depicts the same blot used in Fig. 4A but incubated with an anti-cbl-b specific antibody.

Figure 5A shows an Epidermal Growth Factor (EGF)-induced association of cbl-SL with the EGF Receptor (EGF-R) and Grb2 using a western blot. Fig. 5B shows that an association between Cbl-SL and Grb-2 was detected after EGF stimulation, in comparison to Cbl-Grb-2 association which is detectable both before and after EGF stimulation.

Figure 6 depicts the *in vitro* association of cbl-SL binds with the SH3 domains of Fyn, p85, Nck, as well as with Grb-2, in the presence or absence of EGF stimulation.

Figure 7 depicts a kit comprising an agent of the invention (e.g., anti-cbl-SL Abs, cbl-SL epitopes, etc.), a control agent, and instructions for utilizing such agents in diagnostic or therapeutic applications.

Figure 8 shows the enhanced down-regulation of the EGF-receptor (EGFR) upon cbl-SL overexpression.

Brief Description of the Sequences

SEQ ID NO:1 is the nucleotide sequence of the human cbl-SL cDNA.

SEQ ID NO:2 is the predicted amino acid sequence of the translation product of human cbl-SL cDNA (SEQ ID NO:1).

SEQ ID NO:3 is the nucleotide sequence of the largest open reading frame of the human cbl-SL cDNA of SEQ ID NO:1, encoding for the polypeptide of SEQ ID NO:2.

SEQ ID NO:4 is the nucleotide sequence of a human EST with GenBank Acc. no.: AA113289.

SEQ ID NO:5 is the nucleotide sequence of a human EST with GenBank Acc. no.: AA112513.

SEQ ID NO:6 is the nucleotide sequence of a *sus scrofa* cDNA with EMBL Acc. no.: F22931.

SEQ ID NO:7 is the nucleotide sequence of *C. Elegans* sli-1 cDNA with GenBank Acc. no.: X89223.

SEQ ID NO:8 is the nucleotide sequence of human c-cbl cDNA with GenBank Acc. no.: X57110.

SEQ ID NO:9 is the nucleotide sequence of human cbl-b cDNA with GenBank Acc. no.: U26710.

SEQ ID NO:10 is the amino acid sequence of the cbl-SL peptide (aa 445-461 of SEQ ID NO:2) recognized by one of the anti-cbl-SL specific antibodies of the invention.

Detailed Description of the Invention

One aspect of the invention involves the cloning of a cDNA encoding cbl-SL. Cbl-SL according to the invention is an isolated nucleic acid molecule that comprises a nucleic acid molecule of SEQ ID NO:1, and codes for a ~50kd protein that is believed to play a role in the regulation of a cell's growth, differentiation and proliferation. The sequence of the human cbl-SL cDNA is presented as SEQ ID NO:1, and the predicted amino acid sequence of this cDNA's encoded protein product is presented as SEQ ID NO:2. Cbl-SL associated functions are believed to be mediated by cbl-SL's binding to other molecules and polypeptides. "Cbl-SL activity," as used herein, refers to the specific binding of cbl-SL to a

tyrosine kinase. "Tyrosine kinases" include *receptor* tyrosine kinases and *non-receptor* tyrosine kinases. *Receptor* tyrosine kinases include, but are not limited to, EGRF, PDGFR, CSF-1R (c-fms), c-kit, LET-23R, HGFR/SFR (c-met), FGFR, HER2/neu, HER3, HER4, IGF1R, flt3/flk2, flk1, erbB, c-ret, EphA2, TrkB (BDNFR), tek/tie2, stk, flt-1 (VEGFR),
5 RON, TrkA (NGFR), MuSK, VEGFR2, ROR, tie1, etc. *Non-receptor* tyrosine kinases include, but are not limited to, Fyn, Lck, Lyn, Syk/ZAP-70, Src, Yes, Hck, Blk, Yrk, Fgr, Rak, Brk, Csk. Tyrosine kinases may be phosphorylated or non-phosphorylated, and *receptor* tyrosine kinases may be ligand-activated or not activated. Preferably, Cbl-SL binds phosphorylated tyrosine kinases and ligand-activated *receptor* tyrosine kinases. Cbl-SL
10 activity therefore includes cbl-SL binding at least to the intact Epidermal Growth Factor (EGF) receptor with or without EGF stimulation, or to the Grb-2 (adapter protein) SH2 domain only after EGF stimulation. More preferably, Cbl-SL binds phosphorylated tyrosine kinases and decreases the level of phosphorylation of the tyrosine kinase, and/or downregulates (or inhibits) expression of the tyrosine kinase.

15 As used herein, "downregulating expression" refers to inhibiting (i.e., reducing to a detectable extent) replication, transcription, and/or translation of a tyrosine kinase, since inhibition of any of these processes results in the inhibition of expression of the tyrosine kinase polypeptide encoded by the gene. The term also refers to inhibition of post-translational modifications on the tyrosine kinase polypeptide (e.g., in its phosphorylation),
20 since inhibition of such modifications will also prevent proper expression (i.e., expression as in a wild type cell) of the encoded polypeptide. The term also refers to an increase in, or facilitation of, tyrosine kinase degradation (e.g., *via* increased ubiquitination). Tyrosine kinase turnover can be determined using methods well known in the art and elsewhere herein (see under the Examples section). The inhibition of gene expression can be directly
25 determined by detecting a decrease in the level of mRNA for the gene, or the level of protein expression of the gene, using any suitable means known to the art, such as nucleic acid hybridization or antibody detection methods, respectively. Inhibition of gene expression can also be determined indirectly by detecting a change in tyrosine kinase activity as a whole (e.g., control of cell cycle, induction of cell growth and/or proliferation, control of cell
30 differentiation, control of cell migration, phosphorylation of target polypeptides, etc.).

In contrast to cbl (and cbl-b), cbl-SL does not bind to N-terminal SH3 domains of the adapter protein Nck (adapter proteins function by mediating the rapid and specific assembly of multi-protein complexes during the signal transduction), and thus, cbl-SL activity excludes such binding.

As used herein, a subject is a human, non-human primate, cow, horse, pig, sheep, goat, dog, cat or rodent. In all embodiments human cbl-SL and human subjects are preferred.

Analysis of the sequence by comparison to nucleic acid and protein databases show that cbl-SL shares a limited homology (58% at the amino acid level) to c-cbl (SEQ ID NO:8) and cbl-b (SEQ ID NO:9). Limited homology is also shared between cbl-SL and the *C. Elegans* sli-1 cDNA (SEQ ID NO:7).

The invention thus involves in one aspect an isolated cbl-SL polypeptide, the cDNA encoding this polypeptide, functional modifications and variants of the foregoing, useful fragments of the foregoing, as well as diagnostics and therapeutics relating thereto.

As used herein with respect to nucleic acids, the term "isolated" means: (i) amplified *in vitro* by, for example, polymerase chain reaction (PCR); (ii) recombinantly produced by cloning; (iii) purified, as by cleavage and gel separation; or (iv) synthesized by, for example, chemical synthesis. An isolated nucleic acid is one which is readily manipulable by recombinant DNA techniques well known in the art. Thus, a nucleotide sequence contained in a vector in which 5' and 3' restriction sites are known or for which polymerase chain reaction (PCR) primer sequences have been disclosed is considered isolated but a nucleic acid sequence existing in its native state in its natural host is not. An isolated nucleic acid may be substantially purified, but need not be. For example, a nucleic acid that is isolated within a cloning or expression vector is not pure in that it may comprise only a tiny percentage of the material in the cell in which it resides. Such a nucleic acid is isolated, however, as the term is used herein because it is readily manipulable by standard techniques known to those of ordinary skill in the art.

As used herein with respect to polypeptides, the term "isolated" means separated from its native environment in sufficiently pure form so that it can be manipulated or used for any one of the purposes of the invention. Thus, isolated means sufficiently pure to be used (i) to raise and/or isolate antibodies, (ii) as a reagent in an assay, or (iii) for sequencing, etc.

According to the invention, isolated nucleic acid molecules that code for a cbl-SL polypeptide include: (a) nucleic acid molecules which hybridize under stringent conditions to a molecule consisting of a nucleic acid of SEQ ID NO:1 and which code for a cbl-SL polypeptide, (b) deletions, additions and substitutions of (a) which code for a respective cbl-SL polypeptide, (c) nucleic acid molecules that differ from the nucleic acid molecules of (a) or (b) in codon sequence due to the degeneracy of the genetic code, and (d) complements of (a), (b) or (c).

Homologs and alleles of the cbl-SL nucleic acids of the invention can be identified by conventional techniques. Thus, an aspect of the invention is those nucleic acid sequences which code for cbl-SL polypeptides and which hybridize to a nucleic acid molecule consisting of the coding region of SEQ ID NO:1, under stringent conditions. The term "stringent conditions" as used herein refers to parameters with which the art is familiar. Nucleic acid hybridization parameters may be found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. More specifically, stringent conditions, as used herein, refers, for example, to hybridization at 65°C in hybridization buffer (3.5 x SSC, 0.02% Ficoll, 0.02% polyvinyl pyrrolidone, 0.02% Bovine Serum Albumin, 2.5mM NaH₂PO₄(pH7), 0.5% SDS, 2mM EDTA). SSC is 0.15M sodium chloride/0.15M sodium citrate, pH7; SDS is sodium dodecyl sulphate; and EDTA is ethylenediaminetetracetic acid. After hybridization, the membrane upon which the DNA is transferred is washed at 2 x SSC at room temperature and then at 0.1 x SSC/0.1 x SDS at temperatures up to 68°C.

There are other conditions, reagents, and so forth which can be used, and would result in a similar degree of stringency. The skilled artisan will be familiar with such conditions, and thus they are not given here. It will be understood, however, that the skilled artisan will be able to manipulate the conditions in a manner to permit the clear identification of homologs and alleles of cbl-SL nucleic acids of the invention. The skilled artisan also is familiar with the methodology for screening cells and libraries for expression of such molecules which then are routinely isolated, followed by isolation of the pertinent nucleic acid molecule and sequencing.

In general homologs and alleles typically will share at least 40% nucleotide identity and/or at least 50% amino acid identity to SEQ ID NO:1 and SEQ ID NO:2, respectively, in some instances will share at least 50% nucleotide identity and/or at least 65% amino acid identity and in still other instances will share at least 60% nucleotide identity and/or at least 75% amino acid identity. The homology can be calculated using various, publicly available software tools developed by NCBI (Bethesda, Maryland) that can be obtained through the internet (<ftp://ncbi.nlm.nih.gov/pub/>). Exemplary tools include the BLAST system available at <http://www.ncbi.nlm.nih.gov>. Pairwise and ClustalW alignments (BLOSUM30 matrix setting) as well as Kyte-Doolittle hydropathic analysis can be obtained using the MacVector

sequence analysis software (Oxford Molecular Group). Watson-Crick complements of the foregoing nucleic acids also are embraced by the invention.

In screening for cbl-SL related genes, such as homologs and alleles of cbl-SL, a Southern blot may be performed using the foregoing conditions, together with a radioactive probe. After washing the membrane to which the DNA is finally transferred, the membrane can be placed against X-ray film or a phosphorimager plate to detect the radioactive signal.

Given that the expression of the cbl-SL gene is abundant in certain human tissues, and given the teachings herein of a full-length human cbl-SL cDNA clone, other mammalian sequences such as the mouse cDNA clone corresponding to the human cbl-SL gene can be isolated from a cDNA library prepared from one or more of the tissues in which cbl-SL expression is abundant, using standard colony hybridization techniques.

The invention also includes degenerate nucleic acids which include alternative codons to those present in the native materials. For example, serine residues are encoded by the codons TCA, AGT, TCC, TCG, TCT and AGC. Each of the six codons is equivalent for the purposes of encoding a serine residue. Thus, it will be apparent to one of ordinary skill in the art that any of the serine-encoding nucleotide triplets may be employed to direct the protein synthesis apparatus, *in vitro* or *in vivo*, to incorporate a serine residue into an elongating cbl-SL polypeptide. Similarly, nucleotide sequence triplets which encode other amino acid residues include, but are not limited to: CCA, CCC, CCG and CCT (proline codons); CGA, CGC, CGG, CGT, AGA and AGG (arginine codons); ACA, ACC, ACG and ACT (threonine codons); AAC and AAT (asparagine codons); and ATA, ATC and ATT (isoleucine codons). Other amino acid residues may be encoded similarly by multiple nucleotide sequences. Thus, the invention embraces degenerate nucleic acids that differ from the biologically isolated nucleic acids in codon sequence due to the degeneracy of the genetic code.

The invention also provides isolated unique fragments of SEQ ID NO:1 or SEQ ID NO:3 or complements of thereof. A unique fragment is one that is a 'signature' for the larger nucleic acid. For example, the unique fragment is long enough to assure that its precise sequence is not found in molecules within the human genome outside of the cbl-SL nucleic acids defined above (and human alleles). Those of ordinary skill in the art may apply no more than routine procedures to determine if a fragment is unique within the human genome. Unique fragments, however, exclude fragments completely composed of the nucleotide sequences of any of GenBank accession numbers listed in Table I (AA113289, AA112513, F22931*, X89223, X57110, U26710) or other previously published sequences as of the filing date of this application.

5 A fragment which is completely composed of the sequence described in the foregoing GenBank deposits is one which does not include any of the nucleotides unique to the sequences of the invention. Thus, a unique fragment must contain a nucleotide sequence other than the exact sequence of those in GenBank or fragments thereof. The difference may be an addition, deletion or substitution with respect to the GenBank sequence or it may be a sequence wholly separate from the GenBank sequence.

10 Unique fragments can be used as probes in Southern and Northern blot assays to identify such nucleic acids, or can be used in amplification assays such as those employing PCR. As known to those skilled in the art, large probes such as 200, 250, 300 or more nucleotides are preferred for certain uses such as Southern and Northern blots, while smaller fragments will be preferred for uses such as PCR. Unique fragments also can be used to produce fusion proteins for generating antibodies or determining binding of the polypeptide fragments, as demonstrated in the Examples, or for generating immunoassay components. Likewise, unique fragments can be employed to produce nonfused fragments of the cbl-SL polypeptides, useful, for example, in the preparation of antibodies, immunoassays or therapeutic applications. Unique fragments further can be used as antisense molecules to inhibit the expression of cbl-SL nucleic acids and polypeptides respectively.

20 As will be recognized by those skilled in the art, the size of the unique fragment will depend upon its conservancy in the genetic code. Thus, some regions of SEQ ID NO:1 or SEQ ID NO:3 and complements will require longer segments to be unique while others will require only short segments, typically between 12 and 32 nucleotides long (e.g. 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 and 32 bases) or more, up to the entire length of the disclosed sequence. As mentioned above, this disclosure intends to embrace each and every fragment of each sequence, beginning at the first nucleotide, the second nucleotide and so on, up to 8 nucleotides short of the end, and ending anywhere from nucleotide number 8, 9, 10 and so on for each sequence, up to the very last nucleotide, (provided the sequence is unique as described above). Virtually any segment of the region of SEQ ID NO:1 beginning at nucleotide 1 and ending at nucleotide 1547, or SEQ ID NO:3 beginning at nucleotide 1 and ending at nucleotide 1422, or complements thereof, that is 20 or more nucleotides in length will be unique. Those skilled in the art are well versed in methods for selecting such sequences, typically on the basis of the ability of the unique fragment to selectively distinguish the sequence of interest from other sequences in the human genome of the fragment to those on known databases typically is all that is necessary, although *in vitro* confirmatory hybridization and sequencing analysis may be performed.

As mentioned above, the invention embraces antisense oligonucleotides that selectively bind to a nucleic acid molecule encoding a cbl-SL polypeptide, to decrease cbl-SL activity. When using antisense preparations of the invention, slow intravenous administration is preferred.

As used herein, the term "antisense oligonucleotide" or "antisense" describes an oligonucleotide that is an oligoribonucleotide, oligodeoxyribonucleotide, modified oligoribonucleotide, or modified oligodeoxyribonucleotide which hybridizes under physiological conditions to DNA comprising a particular gene or to an mRNA transcript of that gene and, thereby, inhibits the transcription of that gene and/or the translation of that mRNA. The antisense molecules are designed so as to interfere with transcription or translation of a target gene upon hybridization with the target gene or transcript. Those skilled in the art will recognize that the exact length of the antisense oligonucleotide and its degree of complementarity with its target will depend upon the specific target selected, including the sequence of the target and the particular bases which comprise that sequence. It is preferred that the antisense oligonucleotide be constructed and arranged so as to bind selectively with the target under physiological conditions, i.e., to hybridize substantially more to the target sequence than to any other sequence in the target cell under physiological conditions. Based upon SEQ ID NO:1 or upon allelic or homologous genomic and/or cDNA sequences, one of skill in the art can easily choose and synthesize any of a number of appropriate antisense molecules for use in accordance with the present invention. In order to be sufficiently selective and potent for inhibition, such antisense oligonucleotides should comprise at least 10 and, more preferably, at least 15 consecutive bases which are complementary to the target, although in certain cases modified oligonucleotides as short as 7 bases in length have been used successfully as antisense oligonucleotides (Wagner et al., *Nat. Med.* 1(11):1116-1118, 1995). Most preferably, the antisense oligonucleotides comprise a complementary sequence of 20-30 bases. Although oligonucleotides may be chosen which are antisense to any region of the gene or mRNA transcripts, in preferred embodiments the antisense oligonucleotides correspond to N-terminal or 5' upstream sites such as translation initiation, transcription initiation or promoter sites. In addition, 3'-untranslated regions may be targeted by antisense oligonucleotides. Targeting to mRNA splicing sites has also been used in the art but may be less preferred if alternative mRNA splicing occurs. In addition, the antisense is targeted, preferably, to sites in which mRNA secondary structure is not expected (see, e.g., Sainio et al., *Cell Mol. Neurobiol.* 14(5):439-457, 1994) and at which proteins are not expected to bind. Finally, although, SEQ ID No:1 discloses a cDNA sequence, one of

ordinary skill in the art may easily derive the genomic DNA corresponding to this sequence. Thus, the present invention also provides for antisense oligonucleotides which are complementary to the genomic DNA corresponding to SEQ ID NO:1. Similarly, antisense to allelic or homologous cbl-SL cDNAs and genomic DNAs are enabled without undue experimentation.

In one set of embodiments, the antisense oligonucleotides of the invention may be composed of "natural" deoxyribonucleotides, ribonucleotides, or any combination thereof. That is, the 5' end of one native nucleotide and the 3' end of another native nucleotide may be covalently linked, as in natural systems, via a phosphodiester internucleoside linkage. These oligonucleotides may be prepared by art recognized methods which may be carried out manually or by an automated synthesizer. They also may be produced recombinantly by vectors.

In preferred embodiments, however, the antisense oligonucleotides of the invention also may include "modified" oligonucleotides. That is, the oligonucleotides may be modified in a number of ways which do not prevent them from hybridizing to their target but which enhance their stability or targeting or which otherwise enhance their therapeutic effectiveness.

The term "modified oligonucleotide" as used herein describes an oligonucleotide in which (1) at least two of its nucleotides are covalently linked via a synthetic internucleoside linkage (i.e., a linkage other than a phosphodiester linkage between the 5' end of one nucleotide and the 3' end of another nucleotide) and/or (2) a chemical group not normally associated with nucleic acids has been covalently attached to the oligonucleotide. Preferred synthetic internucleoside linkages are phosphorothioates, alkylphosphonates, phosphorodithioates, phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamides, carboxymethyl esters and peptides.

The term "modified oligonucleotide" also encompasses oligonucleotides with a covalently modified base and/or sugar. For example, modified oligonucleotides include oligonucleotides having backbone sugars which are covalently attached to low molecular weight organic groups other than a hydroxyl group at the 3' position and other than a phosphate group at the 5' position. Thus modified oligonucleotides may include a 2'-O-alkylated ribose group. In addition, modified oligonucleotides may include sugars such as arabinose instead of ribose. The present invention, thus, contemplates pharmaceutical preparations containing modified antisense molecules that are complementary to and hybridizable with, under physiological conditions, nucleic acids encoding cbl-SL polypeptides, together with pharmaceutically acceptable carriers. Antisense oligonucleotides

may be administered as part of a pharmaceutical composition. Such a pharmaceutical composition may include the antisense oligonucleotides in combination with any standard physiologically and/or pharmaceutically acceptable carriers which are known in the art. The compositions should be sterile and contain a therapeutically effective amount of the antisense oligonucleotides in a unit of weight or volume suitable for administration to a patient. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. The term "physiologically acceptable" refers to a non-toxic material that is compatible with a biological system such as a cell, cell culture, tissue, or organism. The characteristics of the carrier will depend on the route of administration. Physiologically and pharmaceutically acceptable carriers include diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials which are well known in the art.

The invention also involves expression vectors coding for cbl-SL proteins and fragments and variants thereof and host cells containing those expression vectors. Virtually any cells, prokaryotic or eukaryotic, which can be transformed with heterologous DNA or RNA and which can be grown or maintained in culture, may be used in the practice of the invention. Examples include bacterial cells such as E.coli and mammalian cells such as mouse, hamster, pig, goat, primate, etc. They may be of a wide variety of tissue types, including mast cells, fibroblasts, oocytes and lymphocytes, and they may be primary cells or cell lines. Specific examples include CHO cells and COS cells. Cell-free transcription systems also may be used in lieu of cells.

As used herein, a "vector" may be any of a number of nucleic acids into which a desired sequence may be inserted by restriction and ligation for transport between different genetic environments or for expression in a host cell. Vectors are typically composed of DNA although RNA vectors are also available. Vectors include, but are not limited to, plasmids, phagemids and virus genomes. A cloning vector is one which is able to replicate in a host cell, and which is further characterized by one or more endonuclease restriction sites at which the vector may be cut in a determinable fashion and into which a desired DNA sequence may be ligated such that the new recombinant vector retains its ability to replicate in the host cell. In the case of plasmids, replication of the desired sequence may occur many times as the plasmid increases in copy number within the host bacterium or just a single time per host before the host reproduces by mitosis. In the case of phage, replication may occur actively during a lytic phase or passively during a lysogenic phase. An expression vector is one into which a desired DNA sequence may be inserted by restriction and ligation such that

it is operably joined to regulatory sequences and may be expressed as an RNA transcript. Vectors may further contain one or more marker sequences suitable for use in the identification of cells which have or have not been transformed or transfected with the vector. Markers include, for example, genes encoding proteins which increase or decrease either resistance or sensitivity to antibiotics or other compounds, genes which encode enzymes whose activities are detectable by standard assays known in the art (e.g., β -galactosidase or alkaline phosphatase), and genes which visibly affect the phenotype of transformed or transfected cells, hosts, colonies or plaques (e.g., green fluorescent protein). Preferred vectors are those capable of autonomous replication and expression of the structural gene products present in the DNA segments to which they are operably joined.

As used herein, a coding sequence and regulatory sequences are said to be "operably" joined when they are covalently linked in such a way as to place the expression or transcription of the coding sequence under the influence or control of the regulatory sequences. If it is desired that the coding sequences be translated into a functional protein, two DNA sequences are said to be operably joined if induction of a promoter in the 5' regulatory sequences results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a promoter region would be operably joined to a coding sequence if the promoter region were capable of effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide.

The precise nature of the regulatory sequences needed for gene expression may vary between species or cell types, but shall in general include, as necessary, 5' non-transcribed and 5' non-translated sequences involved with the initiation of transcription and translation respectively, such as a TATA box, capping sequence, CAAT sequence, and the like. Especially, such 5' non-transcribed regulatory sequences will include a promoter region which includes a promoter sequence for transcriptional control of the operably joined gene. Regulatory sequences may also include enhancer sequences or upstream activator sequences as desired. The vectors of the invention may optionally include 5' leader or signal sequences. The choice and design of an appropriate vector is within the ability and discretion of one of ordinary skill in the art.

Expression vectors containing all the necessary elements for expression are commercially available and known to those skilled in the art. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. Cells are genetically engineered by the introduction into the cells of
5 heterologous DNA (RNA) encoding cbl-SL polypeptide or fragment or variant thereof. That heterologous DNA (RNA) is placed under operable control of transcriptional elements to permit the expression of the heterologous DNA in the host cell.

Preferred systems for mRNA expression in mammalian cells are those such as pRc/CMV (available from Invitrogen, Carlsbad, CA) that contain a selectable marker such as
10 a gene that confers G418 resistance (which facilitates the selection of stably transfected cell lines) and the human cytomegalovirus (CMV) enhancer-promoter sequences. Additionally, suitable for expression in primate or canine cell lines is the pCEP4 vector (Invitrogen, Carlsbad, CA), which contains an Epstein Barr virus (EBV) origin of replication, facilitating the maintenance of plasmid as a multicopy extrachromosomal element. Another expression
15 vector is the pEF-BOS plasmid containing the promoter of polypeptide Elongation Factor 1 α , which stimulates efficiently transcription *in vitro*. The plasmid is described by Mishizuma and Nagata (*Nuc. Acids Res.* 18:5322, 1990), and its use in transfection experiments is disclosed by, for example, Demoulin (*Mol. Cell. Biol.* 16:4710-4716, 1996). Still another preferred expression vector is an adenovirus, described by Stratford-Perricaudet, which is
20 defective for E1 and E3 proteins (*J. Clin. Invest.* 90:626-630, 1992). The use of the adenovirus as an Adeno.P1A recombinant is disclosed by Warnier et al., in intradermal injection in mice for immunization against P1A (*Int. J. Cancer*, 67:303-310, 1996).

The invention also embraces so-called expression kits, which allow the artisan to prepare a desired expression vector or vectors. Such expression kits include at least separate
25 portions of each of the previously discussed coding sequences. Other components may be added, as desired, as long as the previously mentioned sequences, which are required, are included.

It will also be recognized that the invention embraces the use of the above described, cbl-SL cDNA sequence containing expression vectors, to transfect host cells and cell lines,
30 be these prokaryotic (e.g., *E. coli*), or eukaryotic (e.g., CHO cells, COS cells, yeast expression systems and recombinant baculovirus expression in insect cells). Especially useful are mammalian cells such as mouse, hamster, pig, goat, primate, etc. They may be of a wide variety of tissue types, and include primary cells and cell lines. Specific examples include

dendritic cells, U293 cells, peripheral blood leukocytes, bone marrow stem cells and embryonic stem cells. The invention also permits the construction of cbl-SL gene “knock-outs” in cells and in animals, providing materials for studying certain aspects of cbl-SL activity.

5 The invention also provides isolated polypeptides (including whole proteins and partial proteins), encoded by the foregoing cbl-SL nucleic acids, and include the polypeptide of SEQ ID NO:2 and unique fragments thereof. Such polypeptides are useful, for example, alone or as fusion proteins that can associate with a tyrosine kinase to form a complex, and regulate a cell's growth, differentiation and proliferation, to generate antibodies, as
10 components of an immunoassay, etc. Polypeptides can be isolated from biological samples including tissue or cell homogenates, and can also be expressed recombinantly in a variety of prokaryotic and eukaryotic expression systems by constructing an expression vector appropriate to the expression system, introducing the expression vector into the expression system, and isolating the recombinantly expressed protein. Short polypeptides, including
15 antigenic peptides (such as are presented by MHC molecules on the surface of a cell for immune recognition) also can be synthesized chemically using well-established methods of peptide synthesis.

A unique fragment of an cbl-SL polypeptide, in general, has the features and characteristics of unique fragments as discussed above in connection with nucleic acids. As
20 will be recognized by those skilled in the art, the size of the unique fragment will depend upon factors such as whether the fragment constitutes a portion of a conserved protein domain. Thus, some regions of SEQ ID NO:2 will require longer segments to be unique while others will require only short segments, typically between 5 and 12 amino acids (e.g. 5, 6, 7, 8, 9, 10, 11 and 12 amino acids long or more, including each integer up to the full length,
25 474 amino acids long). Virtually any segment of SEQ ID NO:2, excluding the ones that share identity with it (the polypeptides encoded by the nucleic acids of SEQ ID NOs:4, 5, 6, 7, 8 and 9) that is 9 or more amino acids in length will be unique.

Unique fragments of a polypeptide preferably are those fragments which retain a distinct functional capability of the polypeptide. Functional capabilities which can be retained
30 in a unique fragment of a polypeptide include interaction with antibodies, interaction with other polypeptides or fragments thereof (e.g., a tyrosine kinase phosphatase, other signalling molecules), selective binding to receptors (e.g., the EGF receptor), association with certain adapter molecules (e.g., Grb2), and the non-association with other adapter molecules (e.g., Nck). One important activity is the ability to act as a signature for identifying the

polypeptide. Another is the ability to complex with HLA and to provoke in a human an immune response. Those skilled in the art are well versed in methods for selecting unique amino acid sequences, typically on the basis of the ability of the unique fragment to selectively distinguish the sequence of interest from non-family members. A comparison of the sequence of the fragment to those on known databases typically is all that is necessary.

The invention embraces variants of the cbl-SL polypeptides described above. As used herein, a "variant" of a cbl-SL polypeptide is a polypeptide which contains one or more modification to the primary amino acid sequence of a cbl-SL polypeptide. Modifications which create a cbl-SL polypeptide variant are typically made to the nucleic acid which encodes the cbl-SL polypeptide, and can include deletions, point mutations, truncations, amino acid substitutions and addition of amino acids or non-amino acid moieties to: 1) reduce or eliminate an activity of a cbl-SL polypeptide; 2) enhance a property of a cbl-SL polypeptide, such as protein stability in an expression system or the stability of protein-protein binding; 3) provide a novel activity or property to a cbl-SL polypeptide, such as addition of an antigenic epitope or addition of a detectable moiety; or 4) to provide equivalent or better binding to a cbl-SL polypeptide receptor or other molecule (e.g., heparin). Alternatively, modifications can be made directly to the polypeptide, such as by cleavage, addition of a linker molecule, addition of a detectable moiety, such as biotin, addition of a fatty acid, and the like. Modifications also embrace fusion proteins comprising all or part of the cbl-SL amino acid sequence. One of skill in the art will be familiar with methods for predicting the effect on protein conformation of a change in protein sequence, and can thus "design" a variant cbl-SL polypeptide according to known methods. One example of such a method is described by Dahiyat and Mayo in *Science* 278:82-87, 1997, whereby proteins can be designed *de novo*. The method can be applied to a known protein to vary a only a portion of the polypeptide sequence. By applying the computational methods of Dahiyat and Mayo, specific variants of a cancer associated antigen polypeptide can be proposed and tested to determine whether the variant retains a desired conformation.

Variants can include cbl-SL polypeptides which are modified specifically to alter a feature of the polypeptide unrelated to its physiological activity. For example, cysteine residues can be substituted or deleted to prevent unwanted disulfide linkages. Similarly, certain amino acids can be changed to enhance expression of a cbl-SL polypeptide by eliminating proteolysis by proteases in an expression system (e.g., dibasic amino acid residues in yeast expression systems in which KEX2 protease activity is present).

09434708-110599

Mutations of a nucleic acid which encodes a cbl-SL polypeptide preferably preserve the amino acid reading frame of the coding sequence, and preferably do not create regions in the nucleic acid which are likely to hybridize to form secondary structures, such a hairpins or loops, which can be deleterious to expression of the variant polypeptide.

5 Mutations can be made by selecting an amino acid substitution, or by random mutagenesis of a selected site in a nucleic acid which encodes the polypeptide. Variant polypeptides are then expressed and tested for one or more activities to determine which mutation provides a variant polypeptide with the desired properties. Further mutations can be made to variants (or to non-variant cbl-SL polypeptides) which are silent as to the amino acid
10 sequence of the polypeptide, but which provide preferred codons for translation in a particular host. The preferred codons for translation of a nucleic acid in, e.g., E. coli, are well known to those of ordinary skill in the art. Still other mutations can be made to the noncoding sequences of a cbl-SL gene or cDNA clone to enhance expression of the polypeptide.

The skilled artisan will realize that conservative amino acid substitutions may be made
15 in cbl-SL polypeptides to provide functionally equivalent variants of the foregoing polypeptides, i.e., the variants retain the functional capabilities of the cbl-SL polypeptides. As used herein, a "conservative amino acid substitution" refers to an amino acid substitution which does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Variants can be prepared according to methods for altering
20 polypeptide sequence known to one of ordinary skill in the art such as are found in references which compile such methods, e.g. *Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York. Exemplary functionally equivalent variants of the cbl-SL
25 polypeptides include conservative amino acid substitutions of SEQ ID NO:2. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

Thus functionally equivalent variants of cbl-SL polypeptides, i.e., variants of cbl-SL
30 polypeptides which retain the function of the natural cbl-SL polypeptides, are contemplated by the invention. Conservative amino-acid substitutions in the amino acid sequence of cbl-SL polypeptides to produce functionally equivalent variants of cbl-SL polypeptides typically are made by alteration of a nucleic acid encoding cbl-SL polypeptides (SEQ ID NOs:1, 3). Such substitutions can be made by a variety of methods known to one of ordinary skill in the art.

For example, amino acid substitutions may be made by PCR-directed mutation, site-directed mutagenesis according to the method of Kunkel (Kunkel, *Proc. Nat. Acad. Sci. U.S.A.* 82: 488-492, 1985), or by chemical synthesis of a gene encoding a cbl-SL polypeptide. The activity of functionally equivalent fragments of cbl-SL polypeptides can be tested by cloning
5 the gene encoding the altered cbl-SL polypeptide into a bacterial or mammalian expression vector, introducing the vector into an appropriate host cell, expressing the altered cbl-SL polypeptide, and testing for a functional capability of the cbl-SL polypeptides as disclosed herein (e.g., selective binding to the EGF receptor, Grb-2, etc.).

The invention as described herein has a number of uses, some of which are described
10 elsewhere herein. First, the invention permits isolation of cbl-SL polypeptides. A variety of methodologies well-known to the skilled practitioner can be utilized to obtain isolated cbl-SL molecules. The polypeptide may be purified from cells which naturally produce the polypeptide by chromatographic means or immunological recognition. Alternatively, an expression vector may be introduced into cells to cause production of the polypeptide. In
15 another method, mRNA transcripts may be microinjected or otherwise introduced into cells to cause production of the encoded polypeptide. Translation of cbl-SL mRNA in cell-free extracts such as the reticulocyte lysate system also may be used to produce cbl-SL polypeptides. Those skilled in the art also can readily follow known methods for isolating cbl-SL polypeptides. These include, but are not limited to, immunochromatography, HPLC,
20 size-exclusion chromatography, ion-exchange chromatography and immune-affinity chromatography.

The invention also provides, in certain embodiments, "dominant negative" polypeptides derived from cbl-SL polypeptides. A dominant negative polypeptide is an inactive variant of a protein, which, by interacting with the cellular machinery, displaces an
25 active protein from its interaction with the cellular machinery or competes with the active protein, thereby reducing the effect of the active protein. For example, a dominant negative receptor which binds a ligand but does not transmit a signal in response to binding of the ligand can reduce the biological effect of expression of the ligand. Likewise, a dominant negative catalytically-inactive kinase which interacts normally with target proteins but does
30 not phosphorylate the target proteins can reduce phosphorylation of the target proteins in response to a cellular signal. Similarly, a dominant negative transcription factor which binds to a promoter site in the control region of a gene but does not increase gene transcription can reduce the effect of a normal transcription factor by occupying promoter binding sites without increasing transcription.

The end result of the expression of a dominant negative polypeptide in a cell is a reduction in function of active proteins. One of ordinary skill in the art can assess the potential for a dominant negative variant of a protein, and using standard mutagenesis techniques to create one or more dominant negative variant polypeptides. See, e.g., U.S. Patent No. 5,580,723 and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. The skilled artisan then can test the population of mutagenized polypeptides for diminution in a selected and/or for retention of such an activity. Other similar methods for creating and testing dominant negative variants of a protein will be apparent to one of ordinary skill in the art.

The isolation of the cbl-SL cDNA also makes it possible for the artisan to diagnose a disorder characterized by an aberrant expression of cbl-SL. These methods involve determining expression of the cbl-SL gene, and/or cbl-SL polypeptides derived therefrom. In the former situation, such determinations can be carried out via any standard nucleic acid determination assay, including the polymerase chain reaction, or assaying with labeled hybridization probes as exemplified below. In the latter situation, such determination can be carried out via any standard immunological assay using, for example, antibodies which bind to the secreted cbl-SL protein.

The invention also embraces isolated peptide binding agents which, for example, can be antibodies or fragments of antibodies ("binding polypeptides"), having the ability to selectively bind to cbl-SL polypeptides (e.g., SEQ ID NOs: 2 and 10). Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology. In certain embodiments, the invention excludes binding agents (e.g., antibodies) that bind to the polypeptides encoded for by the nucleic acids of SEQ ID NOs: 4, 5, 6, 7, 8 and 9.

Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W.R. (1986) The Experimental Foundations of Modern Immunology Wiley & Sons, Inc., New York; Roitt, I. (1991) Essential Immunology, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F(ab')₂ fragment, retains both of the antigen binding sites of an intact antibody. Similarly, an antibody from which the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment, retains one of the antigen binding sites of an intact antibody molecule. Proceeding further, Fab fragments consist of a

covalently bound antibody light chain and a portion of the antibody heavy chain denoted Fd. The Fd fragments are the major determinant of antibody specificity (a single Fd fragment may be associated with up to ten different light chains without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

5 Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the epitope of the antigen, and framework regions (FRs), which maintain the tertiary structure of the paratope (see, in general, Clark, 1986; Roitt, 1991). In both the heavy chain Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) 10 separated respectively by three complementarity determining regions (CDR1 through CDR3). The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely responsible for antibody specificity.

It is now well-established in the art that the non-CDR regions of a mammalian antibody may be replaced with similar regions of conspecific or heterospecific antibodies 15 while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently joined to human FR and/or Fc/pFc' regions to produce a functional antibody. Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of humanized murine RSV antibodies in which at least a portion of the 20 murine FR regions have been replaced by FR regions of human origin. Such antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

Thus, as will be apparent to one of ordinary skill in the art, the present invention also provides for F(ab')₂, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or 25 FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric F(ab')₂ fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric Fab fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by 30 homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or CDR2 regions have been replaced by homologous human or non-human sequences. The present invention also includes so-called single chain antibodies.

Thus, the invention involves polypeptides of numerous size and type that bind specifically to cbl-SL polypeptides, and complexes of both cbl-SL polypeptides and their

binding partners. These polypeptides may be derived also from sources other than antibody technology. For example, such polypeptide binding agents can be provided by degenerate peptide libraries which can be readily prepared in solution, in immobilized form, as bacterial flagella peptide display libraries or as phage display libraries. Combinatorial libraries also
5 can be synthesized of peptides containing one or more amino acids. Libraries further can be synthesized of peptides and non-peptide synthetic moieties.

Phage display can be particularly effective in identifying binding peptides useful according to the invention. Briefly, one prepares a phage library (using e.g. m13, fd, or lambda phage), displaying inserts from 4 to about 80 amino acid residues using conventional
10 procedures. The inserts may represent, for example, a completely degenerate or biased array. One then can select phage-bearing inserts which bind to the cbl-SL polypeptide or a complex of cbl-SL and a binding partner. This process can be repeated through several cycles of reselection of phage that bind to the cbl-SL polypeptide or complex. Repeated rounds lead to enrichment of phage bearing particular sequences. DNA sequence analysis can be conducted
15 to identify the sequences of the expressed polypeptides. The minimal linear portion of the sequence that binds to the cbl-SL polypeptide or complex can be determined. One can repeat the procedure using a biased library containing inserts containing part or all of the minimal linear portion plus one or more additional degenerate residues upstream or downstream thereof. Yeast two-hybrid screening methods also may be used to identify polypeptides that
20 bind to the cbl-SL polypeptides. Thus, the cbl-SL polypeptides of the invention, or a fragment thereof, or complexes of cbl-SL and a binding partner can be used to screen peptide libraries, including phage display libraries, to identify and select peptide binding partners of the cbl-SL polypeptides of the invention. Such molecules can be used, as described, for screening assays, for purification protocols, for interfering directly with the functioning of
25 cbl-SL and for other purposes that will be apparent to those of ordinary skill in the art.

A cbl-SL polypeptide, or a fragment thereof, also can be used to isolate their native binding partners. Isolation of binding partners may be performed according to well-known methods. For example, isolated cbl-SL polypeptides (that include cbl-SL phosphorylated polypeptides) can be attached to a substrate, and then a solution suspected of containing an
30 cbl-SL binding partner may be applied to the substrate. If the binding partner for cbl-SL polypeptides is present in the solution, then it will bind to the substrate-bound cbl-SL polypeptide. The binding partner then may be isolated. Other proteins which are binding partners for cbl-SL, may be isolated by similar methods without undue experimentation.

The invention also provides novel kits which could be used to measure the levels of the nucleic acids of the invention, expression products of the invention or anti-cbl-SL antibodies. In the case of nucleic acid detection, pairs of primers for amplifying cbl-SL nucleic acids can be included. The preferred kits would include controls such as known amounts of nucleic acid probes, cbl-SL epitopes (such as cbl-SL expression products) or anti-cbl-SL antibodies, as well as instructions or other printed material. In certain embodiments the printed material can characterize risk of developing a cancer based upon the outcome of the assay. The reagents may be packaged in containers and/or coated on wells in predetermined amounts, and the kits may include standard materials such as labeled immunological reagents (such as labeled anti-IgG antibodies) and the like. One kit is a packaged polystyrene microtiter plate coated with cbl-SL protein and a container containing labeled anti-human IgG antibodies. A well of the plate is contacted with, for example, serum, washed and then contacted with the anti-IgG antibody. The label is then detected. A kit embodying features of the present invention, generally designated by the numeral 11, is illustrated in Figure 7. Kit 11 is comprised of the following major elements: packaging 15, an agent of the invention 17, a control agent 19 and instructions 21. Packaging 15 is a box-like structure for holding a vial (or number of vials) containing an agent of the invention 17, a vial (or number of vials) containing a control agent 19, and instructions 21. Individuals skilled in the art can readily modify packaging 15 to suit individual needs.

The invention also embraces a method of screening for the presence of a carcinoma in a subject suspected of having a carcinoma. The method involves (a) characterizing cbl-SL nucleic acid sequences in a test sample, wherein the test sample is obtained from a tissue of the subject, and (b) comparing the cbl-SL nucleic acid sequences of the test sample to cbl-SL nucleic acid sequences of a control sample (wild-type or mutant). An observed alteration or match in a cbl-SL nucleic acid sequence in the test sample as compared to cbl-SL nucleic acid sequences in the control sample, is indicative of the presence of carcinoma in the subject.

By observed "alteration", it is meant that a cbl-SL nucleic acid sequence in the test sample is compared to a control sequence in the control sample, with the control sequence being a wild-type cbl-SL nucleic acid according to the invention (e.g., SEQ ID NO:1 or 3), and the test sample cbl-SL sequence is different to the control sample cbl-SL sequence. Alternatively, an observed "match" may be detected when a cbl-SL nucleic acid sequence in the test sample is compared to a control sequence in the control sample, with the control sequence being a mutant cbl-SL nucleic acid, and the test sample cbl-SL sequence is found identical to the control sample cbl-SL sequence. Appropriate differences and/or matches in

the sequences can be determined with no more than routine experimentation by those of ordinary skill in the art.

The invention also provides methods to measure the level of cbl-SL expression in a subject. This can be performed by first obtaining a test sample from the subject. The test sample can be tissue or biological fluid. Tissues include brain, heart, serum, breast, colon, bladder, uterus, prostate, stomach, testis, ovary, pancreas, pituitary gland, adrenal gland, thyroid gland, salivary gland, mammary gland, kidney, liver, intestine, spleen, thymus, bone marrow, trachea, and lung. In certain embodiments, test samples originate from colon, breast and prostate tissues, and biological fluids include blood, saliva and urine. Both invasive and non-invasive techniques can be used to obtain such samples and are well documented in the art. At the molecular level both PCR and Northern blotting can be used to determine the level of cbl-SL mRNA using products of this invention described earlier, and protocols well known in the art that are found in references which compile such methods. At the protein level, cbl-SL expression can be determined using either polyclonal or monoclonal anti-cbl-SL sera in combination with standard immunological assays. The preferred methods will compare the measured level of cbl-SL expression of the test sample to a control. A control can include a known amount of a nucleic acid probe, a cbl-SL epitope (such as a cbl-SL expression product), or a similar test sample of a subject with a control or 'normal' level of cbl-SL expression.

The invention also embraces a method for treating subjects expressing a mutant cbl-SL. It involves first determining whether the subject, and in particular a specific tissue of the subject, expresses a mutant cbl-SL or a wild-type cbl-SL. As used herein, "wild-type" refers generally to a molecule which is ordinary, common, without defect or affect, and not mutant. An ordinary molecule, also refers generally to sequences or structures that, while they may vary from a canonical sequence or structure, comprise neutral polymorphisms and do not vary in function from a molecule having a non-mutant sequence or structure. According to the invention, a wild-type cbl-SL is, for example, a nucleic acid of SEQ ID NO:1 and its encoded polypeptide presented as SEQ ID NO:2). Wild-type cbl-SL is capable of binding to, for example, the intact Epidermal Growth Factor (EGF, a.k.a. Grb-2) receptor without EGF stimulation, or to the Grb-2 SH2 domain only after EGF stimulation, or to a tyrosine kinase phosphatase, and does not bind to N-terminal SH3 domains of Nck. Conversely, a "mutant" cbl-SL typically has undergone a nucleic acid substitution that results in a non-conservative amino acid substitution at the polypeptide level that changes the cbl-SL's binding characteristics, thus rendering, for example, a proliferative phenotype to the cell. If the tissue

of the subject expresses a mutant cbl-SL, then an effective amount of a wild-type cbl-SL may be administered to the subject in order to increase expression of wild-type cbl-SL in the affected tissue of the subject. Preferred tissues that may benefit from the foregoing therapeutic regimen include tissues with predominant cbl-SL expression, i.e., non-hematopoietic organs, particularly those with a large epithelial component.

The mode of administration and dosage of the agent will vary with the particular stage of the condition being treated, the age and physical condition of the subject being treated, the duration of the treatment, the nature of the concurrent therapy (if any), the specific route of administration, and the like factors within the knowledge and expertise of the health practitioner. In preferred embodiments of the invention cancers expressing a mutant cbl-SL include: biliary tract cancer; brain cancer, including glioblastomas and medulloblastomas; breast cancer; cervical cancer; choriocarcinoma; colon cancer; endometrial cancer; esophageal cancer; gastric cancer; hematological neoplasms, including acute lymphocytic and myelogenous leukemia; multiple myeloma; AIDS associated leukemias and adult T-cell leukemia lymphoma; intraepithelial neoplasms, including Bowen's disease and Paget's disease; liver cancer; lung cancer; lymphomas, including Hodgkin's disease and lymphocytic lymphomas; neuroblastomas; oral cancer, including squamous cell carcinoma; ovarian cancer, including those arising from epithelial cells, stromal cells, germ cells and mesenchymal cells; pancreas cancer; prostate cancer; colorectal cancer; sarcomas, including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma and osteosarcoma; skin cancer, including melanoma, Kaposi's sarcoma, basocellular cancer and squamous cell cancer; testicular cancer, including germinal tumors (seminoma, non-seminoma[teratomas, choriocarcinomas]), stromal tumors and germ cell tumors; thyroid cancer, including thyroid adenocarcinoma and medullar carcinoma; and renal cancer including adenocarcinoma and Wilms tumor. In other embodiments cancers expressing cbl-SL are cancers aberrantly expressing cbl-SL. Aberrant expression is expression of a mutant cbl-SL.

In certain embodiments the agent can be administered, as mentioned earlier, in combination with other anti-cancer agents such as: Acivicin; Aclarubicin; Acodazole Hydrochloride; Acronine; Adozelesin; Aldesleukin; Altretamine; Ambomycin; Ametantrone Acetate; Aminoglutethimide; Amsacrine; Anastrozole; Anthramycin; Asparaginase; Asperlin; Azacitidine; Azetepa; Azotomycin; Batimastat; Benzodepa; Bicalutamide; Bisantrone Hydrochloride; Bisnafide Dimesylate; Bizelesin; Bleomycin Sulfate; Brequinar Sodium; Bropirimine; Busulfan; Cactinomycin; Calusterone; Caracemide; Carbetimer; Carboplatin; Carmustine; Carubicin Hydrochloride; Carzelesin; Cedefingol; Chlorambucil; Cirolemycin;

Cisplatin; Cladribine; Crisnatol Mesylate; Cyclophosphamide; Cytarabine; Dacarbazine;
 Dactinomycin; Daunorubicin Hydrochloride; Decitabine; Dexormaplatin; Dezaguanine;
 Dezaguanine Mesylate; Diaziquone; Docetaxel; Doxorubicin; Doxorubicin Hydrochloride;
 Droloxifene; Droloxifene Citrate; Dromostanolone Propionate; Duazomycin; Edatrexate;
 5 Eflornithine Hydrochloride; Elsamitrucin; Enloplatin; Enpromate; Epipropidine; Epirubicin
 Hydrochloride; Erbulozole; Esorubicin Hydrochloride; Estramustine; Estramustine Phosphate
 Sodium; Etanidazole; Etoposide; Etoposide Phosphate; Etoprine; Fadrozole Hydrochloride;
 Fazarabine; Fenretinide; Floxuridine; Fludarabine Phosphate; Fluorouracil; Flurocitabine;
 Fosquidone; Fostriecin Sodium; Gemcitabine; Gemcitabine Hydrochloride; Hydroxyurea;
 10 Idarubicin Hydrochloride; Ifosfamide; Ilmofofosine; Interferon Alfa-2a; Interferon Alfa-2b;
 Interferon Alfa-n1; Interferon Alfa-n3; Interferon Beta- I a; Interferon Gamma- I b;
 Iproplatin; Irinotecan Hydrochloride; Lanreotide Acetate; Letrozole; Leuprolide Acetate;
 Liarozole Hydrochloride; Lometrexol Sodium; Lomustine; Losoxantrone Hydrochloride;
 Masoprocol; Maytansine; Mechlorethamine Hydrochloride; Megestrol Acetate; Melengestrol
 15 Acetate; Melphalan; Menogaril; Mercaptopurine; Methotrexate; Methotrexate Sodium;
 Metoprine; Meturedapa; Mitindomide; Mitocarcin; Mitocromin; Mitogillin; Mitomalcin;
 Mitomycin; Mitosper; Mitotane; Mitoxantrone Hydrochloride; Mycophenolic Acid;
 Nocodazole; Nogalamycin; Ormaplatin; Oxisuran; Paclitaxel; Pegaspargase; Peliomycin;
 Pentamustine; Peplomycin Sulfate; Perfosfamide; Pipobroman; Pipo sulfan; Piroxantrone
 20 Hydrochloride; Plicamycin; Plomestane; Porfimer Sodium; Porfiromycin; Prednimustine;
 Procarbazine Hydrochloride; Puromycin; Puromycin Hydrochloride; Pyrazofurin; Riboprine;
 Rogletimide; Safingol; Safingol Hydrochloride; Semustine; Simtrazene; Sparfosate Sodium;
 Sparsomycin; Spirogermanium Hydrochloride; Spiromustine; Spiroplatin; Streptonigrin;
 Streptozocin; Sulofenur; Talisomycin; Tecogalan Sodium; Tegafur; Teloxantrone
 25 Hydrochloride; Taxol; Taxotere; Temoporfin; Teniposide; Teroxirone; Testolactone;
 Thiamiprine; Thioguanine; Thiotepa; Tiazofurin; Tirapazamine; Topotecan Hydrochloride;
 Toremifene Citrate; Trestolone Acetate; Triciribine Phosphate; Trimetrexate; Trimetrexate
 Glucuronate; Triptorelin; Tubulozole Hydrochloride; Uracil Mustard; Uredapa; Vapreotide;
 Verteporfin; Vinblastine Sulfate; Vincristine Sulfate; Vindesine; Vindesine Sulfate;
 30 Vinepidine Sulfate; Vinglycinate Sulfate; Vinleurosine Sulfate; Vinorelbine Tartrate;
 Vinrosidine Sulfate; Vinzolidine Sulfate; Vorozole; Zeniplatin; Zinostatin; Zorubicin
 Hydrochloride.

The invention also embraces methods for inhibiting cell proliferation by contacting a
 cell expressing a mutant cbl-SL with the isolated nucleic acids of the invention, or expression

products thereof, in an amount effective to increase cbl-SL activity in the cell and restore proper (i.e., to wild-type standards) cell cycle regulation. Such methods can be very useful in regulating cancerous cell proliferation the both *in vivo* and *in vitro*. The preferred agent is a wild-type cbl-SL or an active fragment thereof.

5 The pharmaceutical preparations, as described above, are administered in effective amounts. The effective amount will depend upon the mode of administration, the particular condition being treated and the desired outcome. It will also depend upon, as discussed above, the stage of the condition, the age and physical condition of the subject, the nature of concurrent therapy, if any, and like factors well known to the medical practitioner. For
10 therapeutic applications, it is that amount sufficient to achieve a medically desirable result. In some cases this is a decrease in cell proliferation.

Generally, doses of active compounds of the present invention would be from about 0.01 mg/kg per day to 1000 mg/kg per day., It is expected that doses ranging from 50-500 mg/kg will be suitable. A variety of administration routes are available. The methods of the
15 invention, generally speaking, may be practiced using any mode of administration that is medically acceptable, meaning any mode that produces effective levels of the active compounds without causing clinically unacceptable adverse effects. Such modes of administration include oral, rectal, topical, nasal, interdermal, or parenteral routes. The term “parenteral” includes subcutaneous, intravenous, intramuscular, or infusion. Intravenous or
20 intramuscular routes are not particularly suitable for long-term therapy and prophylaxis. They could, however, be preferred in emergency situations. Oral administration will be preferred for prophylactic treatment because of the convenience to the patient as well as the dosing schedule. When peptides are used therapeutically, in certain embodiments a desirable route of administration is by pulmonary aerosol. Techniques for preparing aerosol delivery systems
25 containing peptides are well known to those of skill in the art. Generally, such systems should utilize components which will not significantly impair the biological properties of the antibodies, such as the paratope binding capacity (see, for example, Sciarra and Cutie, “Aerosols,” in Remington’s Pharmaceutical Sciences, 18th edition, 1990, pp 1694-1712; incorporated by reference). Those of skill in the art can readily determine the various
30 parameters and conditions for producing antibody or peptide aerosols without resort to undue experimentation.

Compositions suitable for oral administration may be presented as discrete units, such as capsules, tablets, lozenges, each containing a predetermined amount of the active agent.

Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir or an emulsion.

Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like. Lower doses will result from other forms of administration, such as intravenous administration. In the event that a response in a subject is insufficient at the initial doses applied, higher doses (or effectively higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits. Multiple doses per day are contemplated to achieve appropriate systemic levels of compounds.

The cbl-SL polypeptides or fragments thereof may be combined, optionally, with a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid or liquid filler, diluents or encapsulating substances which are suitable for administration into a human. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being co-mingled with the molecules of the present invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficacy.

When administered, the pharmaceutical preparations of the invention are applied in pharmaceutically-acceptable amounts and in pharmaceutically-acceptably compositions. Such preparations may routinely contain salt, buffering agents, preservatives, compatible carriers, and optionally other therapeutic agents. When used in medicine, the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically-acceptable salts thereof and are not excluded from the scope of the invention. Such pharmacologically and pharmaceutically-acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, citric, formic, malonic, succinic, and the

like. Also, pharmaceutically-acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts.

Cbl-SL polypeptides preferably are produced recombinantly, although such polypeptides may be isolated from biological extracts. Recombinantly produced cbl-SL polypeptides include chimeric proteins comprising a fusion of a cbl-SL protein with another polypeptide, e.g., a polypeptide capable of providing or enhancing protein-protein binding, sequence specific nucleic acid binding (such as GAL4), enhancing stability of the cbl-SL polypeptide under assay conditions, or providing a detectable moiety, such as green fluorescent protein. A polypeptide fused to a cbl-SL polypeptide or fragment may also provide means of readily detecting the fusion protein, e.g., by immunological recognition or by fluorescent labeling.

The invention also useful in the generation of transgenic non-human animals. As used herein, "transgenic non-human animals" includes non-human animals having one or more exogenous nucleic acid molecules incorporated in germ line cells and/or somatic cells. Thus the transgenic animal include "knockout" animals having a homozygous or heterozygous gene disruption by homologous recombination, animals having episomal or chromosomally incorporated expression vectors, etc. Knockout animals can be prepared by homologous recombination using embryonic stem cells as is well known in the art. The recombination may be facilitated using, for example, the cre/lox system or other recombinase systems known to one of ordinary skill in the art. In certain embodiments, the recombinase system itself is expressed conditionally, for example, in certain tissues or cell types, at certain embryonic or post-embryonic developmental stages, inducibly by the addition of a compound which increases or decreases expression, and the like. In general, the conditional expression vectors used in such systems use a variety of promoters which confer the desired gene expression pattern (e.g., temporal or spatial). Conditional promoters also can be operably linked to cbl-SL nucleic acid molecules to increase expression of cbl-SL in a regulated or conditional manner. *Trans*-acting negative regulators of cbl-SL activity or expression also can be operably linked to a conditional promoter as described above. Such *trans*-acting regulators include antisense cbl-SL nucleic acids molecules, nucleic acid molecules which encode dominant negative cbl-SL molecules, ribozyme molecules specific for cbl-SL nucleic acids, and the like. The transgenic non-human animals are useful in experiments directed toward testing biochemical or physiological effects of diagnostics or therapeutics for conditions characterized by increased or decreased cbl-SL expression. Other uses will be apparent to one of ordinary skill in the art.

06501-802460

The invention also contemplates gene therapy. The procedure for performing *ex vivo* gene therapy is outlined in U.S. Patent 5,399,346 and in exhibits submitted in the file history of that patent, all of which are publicly available documents. In general, it involves introduction *in vitro* of a functional copy of a gene into a cell(s) of a subject which contains a defective copy of the gene, and returning the genetically engineered cell(s) to the subject. The functional copy of the gene is under operable control of regulatory elements which permit expression of the gene in the genetically engineered cell(s). Numerous transfection and transduction techniques as well as appropriate expression vectors are well known to those of ordinary skill in the art, some of which are described in PCT application WO95/00654. *In vivo* gene therapy using vectors such as adenovirus, retroviruses, herpes virus, and targeted liposomes also is contemplated according to the invention.

The invention also embraces a method for downregulating (or inhibiting) expression of a tyrosine kinase in a cell. The method involves contacting a cell expressing a tyrosine kinase with a cbl-SL polypeptide, in an amount effective to downregulate expression of the tyrosine kinase in the cell. The term "downregulating (or inhibiting) expression" is as described earlier. "Contacting," as used herein, refers to intracellular binding of a cbl-SL polypeptide with a tyrosine kinase. Preferably, cbl-SL polypeptides are produced intracellularly by the cell, following introduction into the cell of cbl-SL nucleic acids of the invention. Cbl-SL polypeptides, however, may be directly introduced into the cell expressing a tyrosine kinase using methods well known in the art, some of which are described elsewhere herein. Preferred cbl-SL polypeptides include the polypeptides of claim 12. In certain embodiments, the tyrosine kinase is selected from the group consisting of a receptor tyrosine kinase, a non-receptor tyrosine kinase, and any of the foregoing tyrosine kinases in complex with another agent. In one embodiment, the tyrosine kinase is ligand activated. In any of the foregoing embodiments, the tyrosine kinase can be phosphorylated. In certain embodiments, the cbl-SL and tyrosine kinase further complexes with an intermediate adapter molecule (e.g., Grb-2).

The invention further provides efficient methods of identifying agents or lead compounds for agents active at the level of a cbl-SL or cbl-SL fragment dependent cellular function. In particular, such functions include interaction with other polypeptides or fragments thereof (e.g., a tyrosine kinase -including selective binding to e.g., the EGF receptor, other signalling molecules, etc.), certain intermediate adapter molecules (e.g., Grb2), and the absence of association with other intermediate adapter molecules (e.g., Nck). Generally, the screening methods involve assaying for compounds which interfere with cbl-

SL activity (such as cbl-SL binding to the EGF receptor), although compounds which enhance cbl-SL activity also can be assayed using the screening methods. Such methods are adaptable to automated, high throughput screening of compounds. The target therapeutic indications for pharmacological agents detected by the screening methods are limited only in that the target cellular function be subject to modulation by alteration of the formation of a complex comprising a cbl-SL polypeptide or fragment thereof and one or more natural cbl-SL binding targets, such as a tyrosine kinase (e.g., EGFR: the EGF receptor, etc). Target indications include cellular processes modulated by cbl-SL (such as the cell cycle, cell migration, cell differentiation, cell proliferation, etc.), and affected by cbl-SL's ability to form complexes with other molecules and polypeptides. The specific binding is typically compared to controls. Controls typically include all of the reagents in the mixture except the candidate agent.

A wide variety of assays for agents are provided, including, labeled *in vitro* protein-protein binding assays, electrophoretic mobility shift assays, immunoassays, cell-based assays such as two- or three-hybrid screens, expression assays, etc. For example, two-hybrid screens are used to rapidly examine the effect of transfected nucleic acids on the intracellular binding of cbl-SL or cbl-SL fragments to specific intracellular targets (e.g. a tyrosine kinase). The transfected nucleic acids can encode, for example, combinatorial peptide libraries or cDNA libraries. Convenient reagents for such assays, e.g., GAL4 fusion proteins, are known in the art. An exemplary cell-based assay involves transfecting a cell with a nucleic acid encoding a cbl-SL polypeptide fused to a GAL4 DNA binding domain and a nucleic acid encoding a reporter gene operably linked to a gene expression regulatory region, such as one or more GAL4 binding sites. Activation of reporter gene transcription occurs when the cbl-SL and reporter fusion polypeptides bind such as to enable transcription of the reporter gene. Agents which modulate a cbl-SL polypeptide mediated cell function are then detected through a change in the expression of reporter gene. Methods for determining changes in the expression of a reporter gene are known in the art.

Cbl-SL fragments used in the methods, when not produced by a transfected nucleic acid are added to an assay mixture as an isolated polypeptide. Cbl-SL polypeptides preferably are produced recombinantly, although such polypeptides may be isolated from biological extracts. Recombinantly produced cbl-SL polypeptides include chimeric proteins comprising a fusion of a cbl-SL protein with another polypeptide, e.g., a polypeptide capable of providing or enhancing protein-protein binding, sequence specific nucleic acid binding (such as GAL4),

enhancing stability of the cbl-SL polypeptide under assay conditions, or providing a detectable moiety, such as green fluorescent protein or Flag epitope.

The assay mixture is comprised, in some embodiments, of a natural intracellular cbl-SL binding target capable of interacting with cbl-SL. While natural cbl-SL binding targets may be used, it is frequently preferred to use portions (e.g., peptides or nucleic acid fragments) or analogs (i.e., agents which mimic the cbl-SL binding properties of the natural binding target for purposes of the assay) of the cbl-SL binding target so long as the portion or analog provides binding affinity and avidity to the cbl-SL fragment measurable in the assay.

The assay mixture also comprises a candidate agent. Typically, a plurality of assay mixtures are run in parallel with different agent concentrations to obtain a different response to the various concentrations. Typically, one of these concentrations serves as a negative control, i.e., at zero concentration of agent or at a concentration of agent below the limits of assay detection. Candidate agents encompass numerous chemical classes, although typically they are organic compounds. Preferably, the candidate pharmacological agents are small organic compounds, i.e., those having a molecular weight of more than 50 yet less than about 2500, preferably less than about 1000 and, more preferably, less than about 500. Candidate agents comprise functional chemical groups necessary for structural interactions with polypeptides and/or nucleic acids, and typically include at least an amine, carbonyl, hydroxyl or carboxyl group, preferably at least two of the functional chemical groups and more preferably at least three of the functional chemical groups. The candidate agents can comprise cyclic carbon or heterocyclic structure and/or aromatic or polyaromatic structures substituted with one or more of the above-identified functional groups. Candidate agents also can be biomolecules such as peptides, saccharides, fatty acids, sterols, isoprenoids, purines, pyrimidines, derivatives or structural analogs of the above, or combinations thereof and the like. Where the agent is a nucleic acid, the agent typically is a DNA or RNA molecule, although modified nucleic acids as defined herein are also contemplated.

Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds and biomolecules, including expression of randomized oligonucleotides, synthetic organic combinatorial libraries, phage display libraries of random peptides, and the like. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural and synthetically produced libraries and compounds can be readily modified through conventional chemical, physical, and biochemical means. Further,

known pharmacological agents may be subjected to directed or random chemical modifications such as acylation, alkylation, esterification, amidification, etc. to produce structural analogs of the agents.

A variety of other reagents also can be included in the mixture. These include reagents such as salts, buffers, neutral proteins (e.g., albumin), detergents, etc. which may be used to facilitate optimal protein-protein and/or protein-nucleic acid binding. Such a reagent may also reduce non-specific or background interactions of the reaction components. Other reagents that improve the efficiency of the assay such as protease, inhibitors, nuclease inhibitors, antimicrobial agents, and the like may also be used.

The mixture of the foregoing assay materials is incubated under conditions whereby, but for the presence of the candidate pharmacological agent, the cbl-SL polypeptide specifically binds the cellular binding target, a portion thereof or analog thereof. The order of addition of components, incubation temperature, time of incubation, and other perimeters of the assay may be readily determined. Such experimentation merely involves optimization of the assay parameters, not the fundamental composition of the assay. Incubation temperatures typically are between 4°C and 40°C. Incubation times preferably are minimized to facilitate rapid, high throughput screening, and typically are between 0.1 and 10 hours.

After incubation, the presence or absence of specific binding between the cbl-SL polypeptide and one or more binding targets is detected by any convenient method available to the user. For cell free binding type assays, a separation step is often used to separate bound from unbound components. The separation step may be accomplished in a variety of ways. Conveniently, at least one of the components is immobilized on a solid substrate, from which the unbound components may be easily separated. The solid substrate can be made of a wide variety of materials and in a wide variety of shapes, e.g., microtiter plate, microbead, dipstick, resin particle, etc. The substrate preferably is chosen to maximum signal to noise ratios, primarily to minimize background binding, as well as for ease of separation and cost.

Separation may be effected for example, by removing a bead or dipstick from a reservoir, emptying or diluting a reservoir such as a microtiter plate well, rinsing a bead, particle, chromatographic column or filter with a wash solution or solvent. The separation step preferably includes multiple rinses or washes. For example, when the solid substrate is a microtiter plate, the wells may be washed several times with a washing solution, which typically includes those components of the incubation mixture that do not participate in specific bindings such as salts, buffer, detergent, non-specific protein, etc. Where the solid

substrate is a magnetic bead, the beads may be washed one or more times with a washing solution and isolated using a magnet.

Detection may be effected in any convenient way for cell-based assays such as two- or three-hybrid screens. The transcript resulting from a reporter gene transcription assay of cbl-SL polypeptide interacting with a target molecule typically encodes a directly or indirectly detectable product, e.g., β -galactosidase activity, luciferase activity, and the like. For cell free binding assays, one of the components usually comprises, or is coupled to, a detectable label. A wide variety of labels can be used, such as those that provide direct detection (e.g., radioactivity, luminescence, optical or electron density, etc). or indirect detection (e.g., epitope tag such as the FLAG epitope, enzyme tag such as horseradish peroxidase, etc.). The label may be bound to a cbl-SL binding partner, or incorporated into the structure of the binding partner.

A variety of methods may be used to detect the label, depending on the nature of the label and other assay components. For example, the label may be detected while bound to the solid substrate or subsequent to separation from the solid substrate. Labels may be directly detected through optical or electron density, radioactive emissions, nonradiative energy transfers, etc. or indirectly detected with antibody conjugates, streptavidin-biotin conjugates, etc. Methods for detecting the labels are well known in the art.

The invention provides cbl-SL-specific binding agents, methods of identifying and making such agents, and their use in diagnosis, therapy and pharmaceutical development. For example, cbl-SL-specific pharmacological agents are useful in a variety of diagnostic and therapeutic applications, especially where disease or disease prognosis is associated with altered cbl-SL binding characteristics. Novel cbl-SL-specific binding agents include cbl-SL-specific antibodies, cell surface receptors, and other natural intracellular and extracellular binding agents identified with assays such as two hybrid screens, and non-natural intracellular and extracellular binding agents identified in screens of chemical libraries and the like.

In general, the specificity of cbl-SL binding to a specific molecule is determined by binding equilibrium constants. Targets which are capable of selectively binding a cbl-SL polypeptide preferably have binding equilibrium constants of at least about 10^7 M^{-1} , more preferably at least about 10^8 M^{-1} , and most preferably at least about 10^9 M^{-1} . The wide variety of cell based and cell free assays may be used to demonstrate cbl-SL-specific binding. Cell based assays include one, two and three hybrid screens, assays in which cbl-SL-mediated transcription is inhibited or increased, etc. Cell free assays include cbl-SL-protein binding

assays, immunoassays, etc. Other assays useful for screening agents which bind cbl-SL polypeptides include fluorescence resonance energy transfer (FRET), and electrophoretic mobility shift analysis (EMSA).

Various techniques may be employed for introducing nucleic acids of the invention into cells, depending on whether the nucleic acids are introduced *in vitro* or *in vivo* in a host. Such techniques include transfection of nucleic acid-CaPO₄ precipitates, transfection of nucleic acids associated with DEAE, transfection with a retrovirus including the nucleic acid of interest, liposome mediated transfection, and the like. For certain uses, it is preferred to target the nucleic acid to particular cells. In such instances, a vehicle used for delivering a nucleic acid of the invention into a cell (e.g., a retrovirus, or other virus; a liposome) can have a targeting molecule attached thereto. For example, a molecule such as an antibody specific for a surface membrane protein on the target cell or a ligand for a receptor on the target cell can be bound to or incorporated within the nucleic acid delivery vehicle. For example, where liposomes are employed to deliver the nucleic acids of the invention, proteins which bind to a surface membrane protein associated with endocytosis may be incorporated into the liposome formulation for targeting and/or to facilitate uptake. Such proteins include capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half life, and the like. Polymeric delivery systems also have been used successfully to deliver nucleic acids into cells, as is known by those skilled in the art. Such systems even permit oral delivery of nucleic acids.

Other delivery systems can include time-release, delayed release or sustained release delivery systems. Such systems can avoid repeated administrations of the anti-inflammatory agent, increasing convenience to the subject and the physician. Many types of release delivery systems are available and known to those of ordinary skill in the art. They include polymer base systems such as poly(lactide-glycolide), copolyoxalates, polycaprolactones, polyesteramides, polyorthoesters, polyhydroxybutyric acid, and polyanhydrides. Microcapsules of the foregoing polymers containing drugs are described in, for example, U.S. Patent 5,075,109. Delivery systems also include non-polymer systems that are: lipids including sterols such as cholesterol, cholesterol esters and fatty acids or neutral fats such as mono- di- and tri-glycerides; hydrogel release systems; sylastic systems; peptide based systems; wax coatings; compressed tablets using conventional binders and excipients; partially fused implants; and the like. Specific examples include, but are not limited to: (a) erosional systems in which an agent of the invention is contained in a form within a matrix

such as those described in U.S. Patent Nos. 4,452,775, 4,675,189, and 5,736,152, and (b) diffusional systems in which an active component permeates at a controlled rate from a polymer such as described in U.S. Patent Nos. 3,854,480, 5,133,974 and 5,407,686. In addition, pump-based hardware delivery systems can be used, some of which are adapted for implantation.

Use of a long-term sustained release implant may be particularly suitable for treatment of chronic conditions. Long-term release, as used herein, means that the implant is constructed and arranged to deliver therapeutic levels of the active ingredient for at least 30 days, and preferably 60 days. Long-term sustained release implants are well-known to those of ordinary skill in the art and include some of the release systems described above.

The invention will be more fully understood by reference to the following examples. These examples, however, are merely intended to illustrate the embodiments of the invention and are not to be construed to limit the scope of the invention.

Examples

Experimental Procedures

Screening of the cDNA library

A pancreatic adenocarcinoma from Stratagene (La Jolla, CA, cat #937208) was screened according to manufacturers recommendation using the insert of I.M.A.G.E. clone ID 526956 from ATCC (Manassas, VA). Hybridization was performed in QuikHyb (Stratagene, La Jolla, CA, cat #201220) under stringent hybridization conditions. Using standard methodology (*Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al., eds., John Wiley & Sons, Inc., New York), individual cDNA clones were subjected to 3-4 rounds of amplification and purification under the same hybridization conditions.

After excision from the vector and subcloning of inserts into the plasmid forms, several clones were sequenced by the Brigham and Women's Hospital Automated Sequencing and Genotyping Facility. Molecular biological techniques such as restriction enzyme treatment, subcloning, DNA extraction, bacterial culture and purification of DNA fragments were performed according to methods well known in the art. Computer analyses of protein and DNA sequences was done using the GCG "Idea" program (Kaneshisa, National Cancer Institute, Frederick, MD, 1986). Restriction endonucleases, expression vectors, and modifying enzymes were purchased from commercial sources (Gibco-BRL). Sequencing

vectors for DNA were purchased from Stratagene (La Jolla, CA). Oligonucleotide probes were purchased from Lofstrand Labs Limited (Gaithersburg, MD). The sequence of the resulting cbl-SL full length clone was confirmed on more independent clones.

Cbl-SL expression

The Tissue blot was purchased from Clontech (Palo Alto, CA; cat #7770-1) and was hybridized using identical hybridization conditions employed in the initial cDNA library screen. The RNA Blot used in the Northern was also a Clontech product (cat #7759-1) that was also subjected to the same hybridization procedures as described above.

Antibodies

The monoclonal antibodies used in this work were 4G10 [anti-Tyr(P)], mouse IgG2a, gift of B. Druker, Oregon Health Science University, Portland, OR) (Druker *et al.*, *N.Engl.J.Med.*, 1989, 321:1383-1391), 12CA5 [anti-influenza hemagglutinin (HA) epitope tag], mouse IgG2b) (Wilson *et al.*, *Cell*, 1984, 37:767-778), and OKT8 (anti-human CD8, mouse IgG2b used as negative control for anti-HA; American Type Culture Collection, Manassas, VA). Polyclonal rabbit antibodies against Grb2 (sc-255), Cbl (sc-170), Cbl-b (sc-1705: cross-reacts with Cbl and Cbl-SL; this paper and data not shown), and EGFR (sc-03) were from Santa Cruz Biotechnology Inc. (Santa Cruz, CA).

Glutathione S-Transferase (GST) Fusion Proteins

Cbl-SL nucleic acid sequences were placed into pGEX4T.1 vector (Pharmacia Biotech, Inc.) in-frame with glutathione S-transferase (GST). Fusion proteins were affinity-purified on glutathione-Sepharose beads (Pharmacia Biotech, Inc.) using art recognized methodology(see e.g., Fukazawa *et al.*, *J. Biol. Chem.*, 1995, 270:19141-19150; Reedquist *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1994, 91:4135-4139; Reedquist *et al.*, *J. Biol. Chem.*, 1996, 271:8435-8442). Multiple independent bacterial colonies producing cbl-SL, c-cbl, Grb-2, Grb-2-SH2, Fyn-SH3, and p85-SH3 fusion proteins were tested. Other/alternative GST-fusion proteins also used in this study have been described previously: GST and GST-Fyn-SH3 in pGEX-2T.K; GST-Src-SH3 and GST-Crk-SH3 in pGEX-2T; GST-Grb2, GST-Grb2-SH2, and GST-p85-SH3 in pGEX-3X (Fukazawa *et al.*, *J. Biol. Chem.*, 1995, 270:19141-19150; Reedquist *et al.*, *J. Biol. Chem.*, 1996, 271:8435-8442). GST-Nck-SH3 (containing N-terminal two SH3 domains of Nck) (Lu *et al.*, *Curr. Biol.*, 1997, 7:85-94) was kindly provided by Bruce Mayer (Children's Hospital, Boston, MA). GST fusion proteins were purified as previously described (Reedquist *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 1994, 91:4135-4139).

Binding Reactions, Immunoprecipitations, Gel Electrophoresis, and Immunoblotting

Binding reactions were typically carried out with 20 µg of purified GST fusion protein and lysate from cells for 1 h (or as indicated) at 4°C. For immunoprecipitations, cell lysate was incubated with antibodies for 1 h (or as indicated) at 4°C, followed by 20 µl of protein A-Sepharose 4B beads (Pharmacia Biotech, Inc., Piscataway, NJ) for 1 h. In all cases, beads were washed six times in lysis buffer (Fukazawa, T. et al., *J Biol Chem*, 1995, 270:19141-19150). Bound proteins were eluted from washed beads in Laemmli sample buffer with 2-mercaptoethanol and resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

Polypeptides were transferred to PVDF membranes (Immunobilon-P, Millipore Corp., Bedford, MA, or polyscreen, NEN-Life Sciences, Inc., Boston, MA), and immunoblotted using protein A-horseradish peroxidase (PA-HRPO) (Cappel-Organon Technika, Durham, NC) as a secondary reagent and enhanced chemiluminescence (ECL) detection (Renaissance™ chemiluminescence reagent and Reflection™ autoradiographic film, NEN-Life Sciences, Inc., Boston, MA), as described in Fukazawa, T. et al., *J Biol Chem*, 1995, 270:19141-19150).

Plasmids, Transient Transfection, EGF Stimulation, and Preparation of Cell Lysates

The full-length Cbl-SL cDNA, encoding an N-terminally hemagglutinin (HA) epitope-tagged Cbl-SL protein, was generated by joining five PCR-derived segments via introduced unique restriction sites which did not change the encoded protein. The HA epitope was placed between residues 1 and 2. The full-length cDNA was cloned into pcDNA3 mammalian expression vector (Invitrogen, Carlsbad, CA) to yield pcDNA3-HA-Cbl-SL. The pAlterMAX-HA-Cbl expression vector, encoding HA-tagged human Cbl, has been described (Lupher *et al.*, *J. Biol. Chem.*, 1996, 271, 24063-24068). The EGFR expression construct was generated by subcloning a *KpnI/SalI* restriction fragment from a pUC-based plasmid (kind gift from Wallace Y. Langdon, University of Western Australia, Nedlands, Western Australia) into the pAlterMAX vector (Lupher, *et al.*, *J. Biol. Chem.*, 1998, 273:35273-35281; Lill *et al.*, *J. Biol. Chem.*, 1999, *in press*).

COS-7 and 293 human embryonic kidney (HEK) cells were grown in DMEM (Life Technologies, Inc., Gaithersburg, MD) supplemented with 10% fetal calf serum (Hyclone Laboratories Inc., Logan, UT), 20 mM HEPES pH 7.35, 1 mM sodium pyruvate, 1 mM non-essential amino acids, 100 units/ml penicillin and 100 µg/ml streptomycin (all from Life Technologies, Inc.). 4.5x10⁵ COS-7 cells per 10-cm dish were plated overnight in DMEM.

The cells were washed and transfected with 2 μ g of pcDNA3-HA-Cbl-SL or p-AlterMAX-HA-Cbl using LipofectamineTM (Life Technologies, Inc.) in OPTI-MEM medium, following the manufacturer's protocol. After 6 hrs, regular DMEM was added. For 293 HEK cells, transient transfection was performed using the calcium phosphate method (Chen & Okayama, *Mol Cell Biol*, 1987, 7:2745-52) using 1 μ g pcDNA3-HA-Cbl-SL or pAlterMAX-HA-Cbl with 0.05 μ g pAlterMAX-EGFR expression plasmids. Culture media was replaced at 14-18h following the addition of DNA precipitates. For EGF stimulation of COS-7 or 293 HEK cells, cells were serum-starved for 5-6h (DMEM containing 0.5% instead of 10% FCS), then were incubated for the indicated lengths of time with purified murine EGF (Sigma) at 100 ng/ml. The cells were then rinsed with ice-cold PBS and processed as described below. Typically, 36-48 hr post-transfection the cells were washed with ice-cold phosphate-buffered saline (PBS) and lysed on ice in a buffer consisting of 0.5% Triton X-100 (Fluka), 50 mM Tris pH 7.5, 150 mM NaCl, 1 mM phenylmethylsulfonyl fluoride, 1 mM sodium orthovanadate, 10 mM sodium fluoride, and a cocktail of protease inhibitors (1 μ g/ml aprotinin, chymostatin, leupeptin, antipain, and pepstatin A) (Sigma Chemicals Co., St Louis, MO). Lysates were rocked at 4°C for 30 min, centrifuged for 10 min and the total protein concentration was determined using the Bradford assay (BioRad Laboratories, Hercules, CA) with bovine serum albumin as the standard. 30 μ g of whole cell lysate was subjected to SDS-PAGE on a 6.5% gel. After transfer, the membrane was cut in two parts just below the 175 kDa marker. The upper part was blotted with anti-EGFR antibody, and the lower one was blotted with a mixture of anti-Cbl antibody and anti-Cbl-SL antiserum (Figure 8). After stripping, both parts were re-probed with the anti-Tyr(P) antibody 4G10; anti-Tyr(P) blot is shown as a single panel (Fig. 8, top panel).

Analysis of Tyrosine Kinase Turnover

The turnover of a tyrosine kinase, e.g., EGF-R in 293 HEK cells expressing endogenous Cbl (mock-transfected) versus those overexpressing HA-Cbl and/or HA-Cbl-SL, was evaluated by pulse-chase analysis using a modification of the previously reported protocol (Lill, et al., *J. Virol.*, 1997, 71:129-137). In brief, cells were transfected in 10cm tissue culture dishes by the calcium phosphate technique, to yield subconfluent cultures at 48-72 h post-transfection. Cells were rinsed with methionine-free DMEM and methionine-starved for 2 h at 37°C by incubation in methionine-free DMEM supplemented with 0.5% dialyzed FBS (Life Technologies, Gaithersburg, MD). Cells were pulse-labeled by adding 50

66901-80242469

μCi/ml EXPRESS labeling mix (NEN/DuPont, Boston, MA) for 40 min at 37°C. After three washes in DMEM, the cells were incubated for 2.5 h in chase medium (DMEM supplemented with 0.5% dialyzed FBS and 3 mg/ml L-methionine) to allow newly synthesized EGF-R to undergo post-translational modification, acquire the capacity to bind to ligand, and be transported to the cell surface. One plate from each transfection set was harvested without further EGF stimulation (0 min stimulation). Cells in the remaining plates were incubated in chase medium containing EGF (100 ng/ml) for the indicated times. Lysate preparation, protein quantification, immunoprecipitation, and gel resolution and PVDF membrane immobilization of proteins were performed as described elsewhere herein. The membrane was used first for autoradiography for detection of radiolabeled proteins (BIOMAX-MR film, Eastman Kodak Co., Rochester, NY), and then in anti-EGF-R immunoblotting to confirm the identity of specific radiolabeled proteins as EGF-R. For quantification of tyrosine kinase (e.g., receptor tyrosine kinase) turnover, autoradiograms were scanned using a Hewlett Packard ScanJet4c™ and Corel Draw™ version 6 software. Densitometry was performed using ScionImage for Windows™ software.

Example 1: Generation of polyclonal rabbit antibodies specific for Cbl-SL

Cos 7 cells were transiently transfected with pcDNA3 (mock), HA-tagged Cbl-SL (in pcDNA3) or HA-tagged Cbl (in pAlterMax). After 48h, cells were lysed in buffer consisting of 50mM Tris pH 8.0, 150 mM NaCl, 0.5% Triton 100X, plus a cocktail of protease and phosphatase inhibitors. 50μg of total lysate was loaded in each lane and resolved on a 7.5% SDS-PAGE gel and transferred to a PDVF membrane. The membrane was blocked in gelatin and blotted with the indicated antibodies for 1h, washed with TBS-Tween and incubated for 30min with Protein A HRP-conjugate. After 1min incubation with a chemiluminescent reagent, autoradiographic exposures were taken. The results of this experiment are shown in Figure 3A. H454 (Santa Cruz) is a polyclonal rabbit anti-Cbl-b antibody which cross-reacts with Cbl (the immunogen corresponds to the N-terminal region of Cbl-b). Anti-Cbl-SL antibody was raised by Zymed Laboratories, Inc. (San Francisco, CA) by immunizing rabbits with cbl-SL peptide (SEQ ID NO:10, or aa 445-461 of SEQ ID NO:2) coupled to KLH. Cbl-SL#2 is the serum from the second rabbit after the first immunization and was used at a dilution 1/1000. Figure 3B depicts the results of an identical experiment, except that two independent anti-Cbl-SL antisera were used. In this second experiment, Cbl-SL#1 and Cbl-SL#2 correspond to the second sera obtained after immunization of the rabbits #1 and #2,

respectively. In Figure 3C, membranes from Fig. 3B were stripped and re-blotted with anti-SL#2 serum in the absence (left membrane) or presence (right membrane) of the immunizing peptide (5µg/ml) as a competitor. Note the absence of Cbl-SL band in the presence of the peptide. In a different experiment, lysates from mock-transfected, Cbl-SL-transfected and Cbl-transfected Cos 7 cells were immunoprecipitated with the anti-HA antibody (12CA5) and the bound proteins were resolved by SDS-PAGE and transferred to a PDVF membrane. The membrane was blotted with the anti-Cbl-SL#2. In Figure 3D, the results of this experiment are presented, showing the specificity of the serum for Cbl-SL only. TL= total lysate; IP-HA= anti-HA immunoprecipitates.

Example 2: Tissue specific expression of Cbl-SL

The Cbl-SL EST (clone # 526956, accession # AA112513) was used to generate a 322 bp ³²P-labelled probe to assess the expression Cbl-SL mRNA in various human tissues (Figure 1B). First, we probed a commercially obtained 'tissue blot' where mRNA samples from various human tissues have been immobilized on a nylon membrane via dot-blotting. The amount of RNA was normalized based on the expression of several housekeeping genes (manufacturer specifications, Clontech). This 'tissue blot' analysis (Figure 1B) showed that Cbl-SL mRNA was predominantly expressed in colon (C4), small intestine (E3), stomach (C8), kidney (E1), liver (E2), pancreas (D3), prostate (C7), mammary gland (D8), salivary gland (D7), trachea (F3), appendix (F1) and placenta (F4), with substantially lower levels in a number of other organs. Most notably, the levels of Cbl-SL mRNA were very low to undetectable in several hematopoietic tissues, in particular the thymus (E5), peripheral blood leukocytes (E6), spleen (E4) and bone marrow (E8). In contrast, Cbl mRNA expression was expectedly broad, with relatively high levels in the hematopoietic tissues (Figure 1A).

Northern blot analysis of selected human tissue RNA samples further confirmed the restricted expression of Cbl-SL (Figure 2), and in addition showed that the predominant mRNA species was about 2.0 kb, compared to a 11 kb size reported for Cbl (Langdon *et al.*, *J.Virol.*, 1989, 63:5420-5424). Minor mRNA species of approximately 4 and 5 kb were observed upon longer exposure of the blot. Collectively, these results demonstrate that the pattern of Cbl-SL mRNA expression is distinct from that of Cbl, and is relatively selective with predominant expression in non-hematopoietic organs, particularly those with a large epithelial component.

Example 3: A candidate endogenous Cbl-SL polypeptide

Different cell lines were lysed and 50µg lysate samples were loaded on a 7.5% SDS-PAGE gel. The membrane was blotted with the anti-Cbl-SL#2 serum. As indicated in Figure 4A, a band detected at about 50kDa co-migrates with the transfected Cbl-SL and corresponds to the endogenous Cbl-SL. The same membrane was stripped and reblotted with the H454 antibody (anti-Cbl-b from Santa Cruz) which reacts with Cbl, Cbl-b and Cbl-SL. The results are depicted in Figure 4B.

Example 4: *Epidermal Growth Factor (EGF)-induced association of Cbl-SL with the EGF Receptor (EGF-R) and Grb2.*

Cos 7 cells were transfected with pcDNA3 (mock), Cbl-SL or Cbl. After 48h, cells were starved for 5 hours in DMEM medium supplemented with 0.5% FCS, and then stimulated for the indicated times with EGF (100 ng/ml). Cells were then lysed with 1ml lysis buffer. 200µg protein samples were immunoprecipitated with anti-HA antibody. After transfer of the gel, the membrane was blotted with a mixture of anti-Cbl-SL + Cbl (Figure 5A, upper blot); after stripping, the membrane was reblotted with anti-pY (Figure 5A, lower blot). Tyrosine phosphorylation of Cbl-SL was not detected in this experiment. However, an associated tyrosine phosphorylated band corresponding to the EGF-R is observed in anti-Cbl-SL immunoprecipitates from lysates of EGF-stimulated cells. In a separate experiment, 400µg of proteins isolated earlier (preceding paragraph), were immunoprecipitated with anti-HA or OKT8 (isotype-matched control) antibody. Immunoprecipitates were separated on a 12% SDS-PAGE gel. After transfer, the membrane was cut in two parts: the upper part was blotted with a mixture of anti-Cbl-SL#2 and anti-Cbl to visualize transfected Cbl-SL and Cbl (as shown in Figure 5B); the lower part (Fig. 5B) was blotted with an anti-Grb-2 antibody (Santa Cruz). In this experiment, association between Cbl-SL and Grb-2 was detected after EGF stimulation, whereas Cbl-Grb-2 association is detectable both before and after EGF stimulation.

Example 5: *In vitro binding of Cbl-SL with GST-fusion proteins of signaling proteins*

Cos 7 cells were transfected with Cbl-SL; after 48h, cells were starved for 5 hours and then stimulated (+) or not (-) with EGF (100 ng/ml) for 5 minutes. 800µg of proteins were mixed for 4 hours with 20µg of different GST-fusion proteins indicated on top of the blot in Figure 6. Precipitates were loaded on a 8% gel. After transfer, the membrane was blotted with a mixture of anti-Cbl-SL#2 and anti-Cbl antibodies. As shown in Figure 6, Cbl-SL binds *in vitro* with the SH3 domains of Fyn and p85, as well as with Grb-2. A very small amount of

Cbl-SL was found to bind to the SH2 domain of Grb-2, only after EGF stimulation. In contrast to Cbl, Cbl-SL did not bind to a GST construct incorporating two N-terminal SH3 domains of Nck (GST-Nck-SH3).

Example 6: *Overexpression of Cbl-SL facilitates the downregulation of the EGFR upon EGF stimulation*

In view of the ability of Cbl-SL to physically interact with EGFR following EGF stimulation in transfected COS-7 cells (see the foregoing examples), we wished to assess if Cbl-SL may play a functional role in EGF signaling pathway. For this purpose, we used a co-transfection system in 293 HEK cells. As 293 HEK cells barely express the EGFR, we co-transfected these cells with an EGFR expression plasmid together with the vector alone, HA-Cbl or HA-Cbl-SL expression plasmids. After 36 hours, cells were serum-starved for 5 hours and stimulated by adding purified murine EGF (100 ng/ml) for the indicated times (Figure 8). The whole cell lysates were then analyzed by anti-Tyr(P) and anti-EGFR immunoblotting.

Overexpression of HA-Cbl (Fig. 8, top panel, lanes 11-15) led to a more rapid decrease in the tyrosine phosphorylation signal on the 175 kDa band corresponding to the EGFR when compared to mock transfectants (lanes 1-5). Concomitantly, a faster decrease in the EGFR protein level was observed (Fig. 8, middle panel). Notably, overexpression of HA-Cbl-SL (lanes 6-10) led to effects comparable to that of HA-Cbl overexpression. For example, significantly lower EGFR protein and phosphotyrosine signals were observed at 20 and 30 minutes after EGF stimulation compared to those in mock transfected cells. Overall, these results indicate that Cbl-SL is capable of negatively regulating the EGFR.

Table I. Sequences with partial homologies to cbl-SL

Sequences with GenBank and EMBL (*) accession numbers:
AA113289, AA112513, F22931*, X89223, X57110, U26710.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims. All references disclosed herein are incorporated by reference in their entirety. What is claimed is presented below and is followed by a Sequence Listing. We claim:

Claims

1. An isolated nucleic acid molecule, comprising:

(a) nucleic acid molecules which hybridize under stringent conditions to a molecule consisting of a nucleic acid of SEQ ID NO:1 and which code for a cbl-SL polypeptide,

(b) deletions, additions and substitutions of (a) which code for a respective cbl-SL polypeptide,

(c) nucleic acid molecules that differ from the nucleic acid molecules of (a) or (b) in codon sequence due to the degeneracy of the genetic code, or

(d) complements of (a), (b) or (c).

2. The isolated nucleic acid molecule of claim 1, wherein the isolated nucleic acid molecule comprises SEQ ID NO:1.

3. The isolated nucleic acid molecule of claim 1, wherein the isolated nucleic acid molecule is the nucleic acid molecule of SEQ ID NO:3 or a fragment thereof.

4. An isolated nucleic acid molecule selected from the group consisting of

(a) a unique fragment of nucleic acid molecule of SEQ ID NO:1,

(b) complements of (a),

provided that the unique fragment includes a sequence of contiguous nucleotides which is not identical to any sequence selected from the sequence group consisting of

(1) sequences having the database accession numbers of Table I,

(2) complements of (1), and

(3) fragments of (1) and (2).

5. The isolated nucleic acid molecule of claim 4, wherein the sequence of contiguous nucleotides is selected from the group consisting of:

(1) at least two contiguous nucleotides nonidentical to the sequence group,

(2) at least three contiguous nucleotides nonidentical to the sequence group,

(3) at least four contiguous nucleotides nonidentical to the sequence group,

(4) at least five contiguous nucleotides nonidentical to the sequence group,

(5) at least six contiguous nucleotides nonidentical to the sequence group,

(6) at least seven contiguous nucleotides nonidentical to the sequence group.

6. The isolated nucleic acid molecule of claim 4, wherein the fragment has a size selected from the group consisting of at least: 8 nucleotides, 10 nucleotides, 12 nucleotides, 14 nucleotides, 16 nucleotides, 18 nucleotides, 20, nucleotides, 22 nucleotides, 24 nucleotides, 26 nucleotides, 28 nucleotides, 30 nucleotides, 50 nucleotides, 75 nucleotides, 100 nucleotides, and 200 nucleotides.
7. The isolated nucleic acid molecule of claim 4, wherein the molecule encodes a polypeptide which is immunogenic.
8. An expression vector comprising the isolated nucleic acid molecule of claims 1, 2, 3, 4, 5, 6, or 7 operably linked to a promoter.
9. An expression vector comprising the isolated nucleic acid molecule of claim 4 operably linked to a promoter.
10. A host cell transformed or transfected with the expression vector of claim 8.
11. A host cell transformed or transfected with the expression vector of claim 9.
12. An isolated polypeptide encoded by the isolated nucleic acid molecule of claim 1, 2, 3, or 4, wherein the polypeptide, or fragment of the polypeptide, has cbl-SL activity.
13. The isolated polypeptide of claim 12, wherein the isolated polypeptide is encoded by the isolated nucleic acid molecule of claim 2.
14. The isolated polypeptide of claim 13, wherein the isolated polypeptide comprises a polypeptide having the sequence of amino acids 1-474 of SEQ ID NO:2.
15. An isolated polypeptide encoded by the isolated nucleic acid molecule of claim 1, 2, or 3, wherein the polypeptide, or fragment of the polypeptide, is immunogenic.
16. The fragment of claim 15, wherein the fragment, or portion of the fragment, binds to a human antibody.
17. An isolated binding polypeptide which binds selectively a polypeptide encoded by the isolated nucleic acid molecule of claim 1, 2, 3, or 4.

18. The isolated binding polypeptide of claim 17, wherein the isolated binding polypeptide binds to a polypeptide having the sequence of amino acids of SEQ ID NO:2 or SEQ ID NO:10.

19. The isolated binding polypeptide of claim 17, wherein the isolated binding polypeptide binds to a polypeptide having the sequence of amino acids of SEQ ID NO:2.

20. The isolated binding polypeptide of claim 19, wherein the isolated binding polypeptide is an antibody or an antibody fragment selected from the group consisting of a Fab fragment, a F(ab)₂ fragment or a fragment including a CDR3 region selective for the polypeptide having the sequence of amino acids of SEQ ID NO:2.

21. An isolated polypeptide comprising a fragment of the polypeptide of claim 12 of sufficient length to represent a sequence unique within the human genome and identifying a polypeptide that has cbl-SL activity, provided that the fragment excludes a sequence of contiguous amino acids encoded by an isolated nucleic acid selected from the group consisting of SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, and SEQ ID NO:9.

22. A kit, comprising a package containing:

an agent that selectively binds to the isolated nucleic acid of claim 1 or an expression product thereof, and

a control for comparing to a measured value of binding of said agent to said isolated nucleic acid of claim 1 or expression product thereof.

23. The kit of claim 22, wherein the control is a predetermined value for comparing to the measured value.

24. The kit of claim 23, wherein the control comprises an epitope of the expression product of the nucleic acid of claim 1.

25. A method for determining the level of cbl-SL expression in a subject, comprising:

- a) obtaining a test sample from a subject,
- b) measuring the expression of cbl-SL in the test sample,
- c) comparing the measured expression of cbl-SL to a control.

26. The method of claim 25, wherein the expression of cbl-SL in (b) is cbl-SL mRNA expression.

27. The method of claim 25, wherein the expression of cbl-SL in (b) is cbl-SL polypeptide expression.

5 28. The method of claim 25, wherein the test sample is tissue.

29. The method of claim 25, wherein the test sample is a biological fluid.

30. The method of claim 26, wherein cbl-SL mRNA expression is measured using the Polymerase Chain Reaction (PCR).

10 31. The method of claim 26, wherein cbl-SL mRNA expression is measured using northern blotting.

32. The method of claim 27, wherein cbl-SL polypeptide expression is measured using monoclonal antisera to cbl-SL.

33. The method of claim 27, wherein cbl-SL polypeptide expression is characterized using polyclonal antisera to cbl-SL.

15 34. A method of screening for the presence of a carcinoma in a subject suspected of having a carcinoma, comprising:

(a) characterizing cbl-SL nucleic acid sequences in a test sample, wherein the test sample is obtained from the subject;

20 (b) comparing the cbl-SL nucleic acid sequences of the test sample to cbl-SL nucleic acid sequences of a control sample,

wherein an observed alteration or match in a cbl-SL nucleic acid sequence in the tissue sample as compared to the cbl-SL nucleic acid sequences in the control sample, is indicative of the presence of carcinoma in the subject.

25 35. The method of claim 34, wherein the observed alteration is apparent when a cbl-SL nucleic acid sequence in the tissue sample is compared to wild-type cbl-SL nucleic acid sequences in the control sample.

36. The method of claim 34, wherein the observed match is apparent when a cbl-SL nucleic acid sequence in the tissue sample is compared to mutant cbl-SL nucleic acid sequences in the control sample.

37. The method of claim 34, wherein cbl-SL mRNA molecules are compared.

38. The method of claim 37, wherein alteration of cbl-SL mRNA is detected by hybridization of mRNA from said tissue sample to a cbl-SL nucleic acid of claim 1, 2, or 3.

39. The method of claim 38, wherein cbl-SL cDNA sequences are compared, said comparing performed by hybridization of a cbl-SL cDNA probe to genomic DNA isolated from said tissue sample.

40. The method of claim 39, further comprising:

(a) subjecting genomic DNA isolated from a non-neoplastic tissue of the subject to Southern hybridization with the cbl-SL cDNA probe; and

(b) comparing the hybridizations of:

(i) the cbl-SL cDNA probe to said tissue sample and (ii) the cbl-SL cDNA probe to said non-neoplastic tissues.

41. The method of claim 39, wherein the cbl-SL cDNA probe detects a restriction fragment length polymorphism.

42. The method of claim 34, wherein cbl-SL nucleic acid sequences are compared, said comparing being performed by determining the sequence of all or part of a cbl-SL cDNA in said tissue sample using a polymerase chain reaction,

wherein deviations in the Cbl-SL cDNA determined from that of the wild-type Cbl-SL nucleic acid sequence shown in SEQ ID NO: 1, is indicative of the presence of the carcinoma in the subject.

43. The method of claim 34, wherein the alteration of cbl-SL nucleic acid sequences is detected by identifying a mismatch between molecules (a) a cbl-SL cDNA or cbl-SL mRNA isolated from said tissue and (b) a nucleic acid probe complementary to the human wild-type

cbl-SL nucleic acid sequence, when molecules (a) and (b) are hybridized to each other to form a duplex.

44. The method of claim 34, wherein cbl-SL nucleic acid sequences are compared and the alteration of cbl-SL nucleic acid sequences is detected by the steps of:

5 (a) amplifying Cbl-SL cDNA sequences in said tissue sample, and

(b) hybridizing the amplified cbl-SL cDNA sequences to nucleic acid probes which comprise Cbl-SL sequences.

10 45. The method of claim 34, wherein cbl-SL nucleic acid sequences are compared and the alteration of Cbl-SL nucleic acid sequences is detected by molecular cloning of Cbl-SL genes in said tissue sample and sequencing all or part of the cloned Cbl-SL gene.

46. The method of claim 34, wherein the detection of alteration of Cbl-SL nucleic acid sequences comprises screening for a deletion mutation.

15 47. The method of claim 34, wherein the detection of alteration of Cbl-SL nucleic acid sequences comprises screening for a point mutation.

48. The method of claim 34, wherein the detection of alteration of Cbl-SL nucleic acid sequences comprises screening for an insertion mutation.

20 49. The method of claim 34, wherein said tissue is selected from the group consisting of brain, heart, serum, breast, colon, bladder, uterus, prostate, stomach, testis, ovary, pancreas, pituitary gland, adrenal gland, thyroid gland, salivary gland, mammary gland, kidney, liver, intestine, spleen, thymus, bone marrow, trachea, and lung.

50. A pharmaceutical composition comprising:

a pharmaceutically effective amount of an agent comprising of an isolated nucleic acid molecule of claim 1 or an expression product thereof, and

25 a pharmaceutically acceptable carrier.

51. The pharmaceutical composition of claim 50, wherein the agent is an expression product of the isolated nucleic acid molecule of claim 1.

52. A method for determining the level of cbl-SL expression in a subject, comprising:

a) measuring the expression of cbl-SL in a test sample, wherein the test sample is obtained from the subject; and

b) comparing the measured expression of cbl-SL to a control.

53. A method for increasing cbl-SL expression in a subject that expresses a mutant cbl-SL, comprising:

administering an isolated cbl-SL nucleic acid molecule of the invention or an expression product thereof to a subject expressing a mutant cbl-SL, in an amount effective to increase wild-type cbl-SL expression in the subject.

54. A method for downregulating expression of a tyrosine kinase in a cell, comprising:

contacting a cell expressing a tyrosine kinase with a cbl-SL polypeptide, in an amount effective to downregulate expression of the tyrosine kinase in the cell.

55. The method of claim 54, wherein the tyrosine kinase is selected from the group consisting of a receptor tyrosine kinase and a non- receptor tyrosine kinase.

56. The method of claim 54, wherein the receptor tyrosine kinase is EGFR.

57. The method of claim 54, wherein the cbl-SL polypeptide is a polypeptide having an amino acid sequence consisting of the sequence set forth in SEQ ID NO:2.

58. The method of claim 54, wherein the cbl-SL polypeptide is encoded by the nucleic acid of SEQ ID NO:1.

59. The method of claim 54, wherein the tyrosine kinase is ligand activated.

60. The method according to anyone of claims 54-58, wherein the tyrosine kinase is phosphorylated.

61. A method for identifying lead compounds for an agent that regulates cbl-SL activity, comprising:

(a) forming a mixture of a cbl-SL polypeptide, a tyrosine kinase that binds a cbl-SL polypeptide, and a candidate agent,

(b) incubating the mixture under conditions which permit specific binding of the tyrosine kinase that binds a cbl-SL polypeptide to the cbl-SL polypeptide,

(c) detecting a reference specific binding of the tyrosine kinase that binds a cbl-SL polypeptide to the cbl-SL polypeptide, the reference specific binding being indicative of cbl-SL activity, and

(d) comparing the reference specific binding to a control.

62. The method of claim 61, wherein the tyrosine kinase is selected from the group consisting of a receptor tyrosine kinase, a non- receptor tyrosine kinase, and any of the foregoing tyrosine kinases in complex with another agent.

63. The method of claim 61, wherein the receptor tyrosine kinase is EGFR.

64. The method of claim 61, wherein the cbl-SL polypeptide is a polypeptide having an amino acid sequence consisting of the sequence set forth in SEQ ID NO:2.

65. The method of claim 61, wherein the cbl-SL polypeptide is encoded by the nucleic acid of SEQ ID NO:1.

66. The method of claim 61, wherein the tyrosine kinase is ligand activated.

67. The method according to anyone of claims 61-65, wherein the tyrosine kinase is phosphorylated.

68. The method of claim 61, further comprising adding an intermediate adapter into the mixture.

69. The method of claim 68, wherein the intermediate adapter is Grb2.

Abstract

The invention pertains to nucleic acids encoding a cbl-SL protein, including fragments and biologically functional variants thereof. The invention also pertains to therapeutics and diagnostics involving the foregoing proteins and genes and agents that bind the foregoing
5 proteins and genes.

09434709.110599

FIGURE 1.




FIGURE 2.




FIGURE 3A.




FIGURE 3B.




FIGURE 3C.




FIGURE 3D.




FIGURE 4A.




FIGURE 4B.




FIGURE 5A.




FIGURE 5B.





FIGURE 6.



05507-80242450

Figure 7.





Figure 8



SEQUENCE LISTING

<110> Borriello, Francescopaolo
 Band, Hamid

<120> Characterization of a novel gene Chl-SL

<130> B0801/7159/ERP

<150> U.S. 60/107,470

<151> 1998-11-06

<160> 10

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1547

<212> DNA

<213> Homo Sapiens

<220>

<221> CDS

<222> (13)...(1434)

<400> 1

cgcgaggctc cc atg gct ctg gcg gtg gcc ccg tgg ggg cga cag tgg gaa	51
Met Ala Leu Ala Val Ala Pro Trp Gly Arg Gln Trp Glu	
1 5 10	
 gag gcc cgc gcc ctg ggc cgg gca gtc agg atg ctg cag cgc cta gaa	99
Glu Ala Arg Ala Leu Gly Arg Ala Val Arg Met Leu Gln Arg Leu Glu	
15 20 25	
 gag caa tgc gtc gac ccc cgg ctg tcc gtg agt ccc cct tcg ctg cgg	147
Glu Gln Cys Val Asp Pro Arg Leu Ser Val Ser Pro Pro Ser Leu Arg	
30 35 40 45	
 gac ctg ctg ccc cgc aca gcg cag ctg ctt cga gag gtg gcc cat tct	195
Asp Leu Leu Pro Arg Thr Ala Gln Leu Leu Arg Glu Val Ala His Ser	
50 55 60	
 cgg cgg gcg gcc ggc gga ggc ggc ccc ggg ggt ccc ggc ggc tct ggg	243
Arg Arg Ala Ala Gly Gly Gly Gly Pro Gly Gly Pro Gly Gly Ser Gly	
65 70 75	
 gac ttt cta ctc atc tac ctg gcc aat ctg gag gcc aag agc agg cag	291
Asp Phe Leu Leu Ile Tyr Leu Ala Asn Leu Glu Ala Lys Ser Arg Gln	
80 85 90	
 gtg gcc gcg ctg ctg cct ccc cgg ggc cga agg agt gcc aac gac gag	339
Val Ala Ala Leu Leu Pro Pro Arg Gly Arg Arg Ser Ala Asn Asp Glu	
95 100 105	

09434708 110599

ctc ttc cgg gcg ggc tcc aga ctc agg cga cag ctg gcc aag ctg gcc 387
 Leu Phe Arg Ala Gly Ser Arg Leu Arg Arg Gln Leu Ala Lys Leu Ala
 110 115 120 125

atc atc ttc agc cac atg cac gca gag ctg cac gca ctc ttc ccc ggg 435
 Ile Ile Phe Ser His Met His Ala Glu Leu His Ala Leu Phe Pro Gly
 130 135 140

gca aag tac tgt gga cac atg tac cag ctc acc aag gcc ccc gcc cac 483
 Ala Lys Tyr Cys Gly His Met Tyr Gln Leu Thr Lys Ala Pro Ala His
 145 150 155

acc ttc tgg agg gaa agt tgc gga gcc cgg tgt gtg ctg ccc tgg gct 531
 Thr Phe Trp Arg Glu Ser Cys Gly Ala Arg Cys Val Leu Pro Trp Ala
 160 165 170

gag ttt gag tcc ctc ctg ggc acc tgc cac cct gtg gaa cca ggc tgc 579
 Glu Phe Glu Ser Leu Leu Gly Thr Cys His Pro Val Glu Pro Gly Cys
 175 180 185

aca gcc ctg gcc ttg cgc acc acc att gac ctc acc tgc agc ggg cac 627
 Thr Ala Leu Ala Leu Arg Thr Thr Ile Asp Leu Thr Cys Ser Gly His
 190 195 200 205

gtg tcc atc ttc gag ttc gac gtc ttc acc agg ctc ttt cag cca tgg 675
 Val Ser Ile Phe Glu Phe Asp Val Phe Thr Arg Leu Phe Gln Pro Trp
 210 215 220

cca aca ctc ctc aag aac tgg cag ctc ctg gca gtc aac cac cca ggc 723
 Pro Thr Leu Leu Lys Asn Trp Gln Leu Leu Ala Val Asn His Pro Gly
 225 230 235

tac atg gcc ttc ctc acc tat gat gag gtc caa gag cgt ctg cag gcc 771
 Tyr Met Ala Phe Leu Thr Tyr Asp Glu Val Gln Glu Arg Leu Gln Ala
 240 245 250

tgc agg gac aag cca ggc agt tac atc ttc cgg ccc agc tgt act cgc 819
 Cys Arg Asp Lys Pro Gly Ser Tyr Ile Phe Arg Pro Ser Cys Thr Arg
 255 260 265

ctg ggg cag tgg gcc atc ggc tat gtg agc tca gat ggc agc atc ctg 867
 Leu Gly Gln Trp Ala Ile Gly Tyr Val Ser Ser Asp Gly Ser Ile Leu
 270 275 280 285

cag acc atc cct gcc aac aaa ccc ctg tcc cag gtg ctc ctg gag gga 915
 Gln Thr Ile Pro Ala Asn Lys Pro Leu Ser Gln Val Leu Leu Glu Gly
 290 295 300

cag aag gac ggc ttc tac ctc tac cca gat gga aag acc cac aac cca 963
 Gln Lys Asp Gly Phe Tyr Leu Tyr Pro Asp Gly Lys Thr His Asn Pro
 305 310 315

gac ctg act gag ctc ggc cag gca gaa ccc cag cag cgc atc cac gtg 1011
 Asp Leu Thr Glu Leu Gly Gln Ala Glu Pro Gln Gln Arg Ile His Val
 320 325 330

tca gag gag cag ctg cag ctc tac tgg gcc atg gac tcc aca ttt gag	1059
Ser Glu Glu Gln Leu Gln Leu Tyr Trp Ala Met Asp Ser Thr Phe Glu	
335 340 345	
ctc tgc aag atc tgt gct gag agc aac aag gat gtg aag att gag ccg	1107
Leu Cys Lys Ile Cys Ala Glu Ser Asn Lys Asp Val Lys Ile Glu Pro	
350 355 360 365	
tgc ggg cac ctg ctc tgc agc tgc tgc ctg gct gcc tgg cag cac tcg	1155
Cys Gly His Leu Leu Cys Ser Cys Cys Leu Ala Ala Trp Gln His Ser	
370 375 380	
gac agc cag acc tgc ccc ttc tgc cgc tgc gag atc aag ggc tgg gag	1203
Asp Ser Gln Thr Cys Pro Phe Cys Arg Cys Glu Ile Lys Gly Trp Glu	
385 390 395	
gcc gtg agt atc tac cag ttc cac ggt cag gct act gct gag gac tca	1251
Ala Val Ser Ile Tyr Gln Phe His Gly Gln Ala Thr Ala Glu Asp Ser	
400 405 410	
ggg aac agc agt gac cag gaa ggc agg gag ttg gag ctg ggg cag gtg	1299
Gly Asn Ser Ser Asp Gln Glu Gly Arg Glu Leu Glu Leu Gly Gln Val	
415 420 425	
ccc ctt tcg gct cct cca ttg ccc cca cgg cca gat ctg ccc ccc agg	1347
Pro Leu Ser Ala Pro Pro Leu Pro Pro Arg Pro Asp Leu Pro Pro Arg	
430 435 440 445	
aag ccc aga aat gcc cag ccg aaa gtg aga ctc cta aag ggg aac tcc	1395
Lys Pro Arg Asn Ala Gln Pro Lys Val Arg Leu Leu Lys Gly Asn Ser	
450 455 460	
cct cca gct gcg ctg gga ccc cag gac cct gcc ccg gcc tgaaggccag	1444
Pro Pro Ala Ala Leu Gly Pro Gln Asp Pro Ala Pro Ala	
465 470	
ggcaccacaga tgtgtgtgtc aagggagccc caagggctgg aaggggggttg tgaaaccgaa	1504
ataaactgcc aagcctgggtc tgtcaaaaaa aaaaaaaaaa aaa	1547

<210> 2
 <211> 474
 <212> PRT
 <213> Homo Sapiens

<400> 2

Met Ala Leu Ala Val Ala Pro Trp Gly Arg Gln Trp Glu Glu Ala Arg	
1 5 10 15	
Ala Leu Gly Arg Ala Val Arg Met Leu Gln Arg Leu Glu Glu Gln Cys	
20 25 30	
Val Asp Pro Arg Leu Ser Val Ser Pro Pro Ser Leu Arg Asp Leu Leu	
35 40 45	
Pro Arg Thr Ala Gln Leu Leu Arg Glu Val Ala His Ser Arg Arg Ala	
50 55 60	
Ala Gly Gly Gly Gly Pro Gly Gly Pro Gly Gly Ser Gly Asp Phe Leu	
65 70 75 80	
Leu Ile Tyr Leu Ala Asn Leu Glu Ala Lys Ser Arg Gln Val Ala Ala	

				85					90				95		
Leu	Leu	Pro	Pro	Arg	Gly	Arg	Arg	Ser	Ala	Asn	Asp	Glu	Leu	Phe	Arg
			100					105					110		
Ala	Gly	Ser	Arg	Leu	Arg	Arg	Gln	Leu	Ala	Lys	Leu	Ala	Ile	Ile	Phe
		115					120					125			
Ser	His	Met	His	Ala	Glu	Leu	His	Ala	Leu	Phe	Pro	Gly	Ala	Lys	Tyr
	130					135					140				
Cys	Gly	His	Met	Tyr	Gln	Leu	Thr	Lys	Ala	Pro	Ala	His	Thr	Phe	Trp
145					150					155					160
Arg	Glu	Ser	Cys	Gly	Ala	Arg	Cys	Val	Leu	Pro	Trp	Ala	Glu	Phe	Glu
			165					170						175	
Ser	Leu	Leu	Gly	Thr	Cys	His	Pro	Val	Glu	Pro	Gly	Cys	Thr	Ala	Leu
			180				185						190		
Ala	Leu	Arg	Thr	Thr	Ile	Asp	Leu	Thr	Cys	Ser	Gly	His	Val	Ser	Ile
		195					200					205			
Phe	Glu	Phe	Asp	Val	Phe	Thr	Arg	Leu	Phe	Gln	Pro	Trp	Pro	Thr	Leu
	210					215					220				
Leu	Lys	Asn	Trp	Gln	Leu	Leu	Ala	Val	Asn	His	Pro	Gly	Tyr	Met	Ala
225				230					235						240
Phe	Leu	Thr	Tyr	Asp	Glu	Val	Gln	Glu	Arg	Leu	Gln	Ala	Cys	Arg	Asp
			245					250					255		
Lys	Pro	Gly	Ser	Tyr	Ile	Phe	Arg	Pro	Ser	Cys	Thr	Arg	Leu	Gly	Gln
		260					265						270		
Trp	Ala	Ile	Gly	Tyr	Val	Ser	Ser	Asp	Gly	Ser	Ile	Leu	Gln	Thr	Ile
		275					280					285			
Pro	Ala	Asn	Lys	Pro	Leu	Ser	Gln	Val	Leu	Leu	Glu	Gly	Gln	Lys	Asp
	290					295					300				
Gly	Phe	Tyr	Leu	Tyr	Pro	Asp	Gly	Lys	Thr	His	Asn	Pro	Asp	Leu	Thr
305				310						315					320
Glu	Leu	Gly	Gln	Ala	Glu	Pro	Gln	Gln	Arg	Ile	His	Val	Ser	Glu	Glu
			325					330						335	
Gln	Leu	Gln	Leu	Tyr	Trp	Ala	Met	Asp	Ser	Thr	Phe	Glu	Leu	Cys	Lys
			340				345						350		
Ile	Cys	Ala	Glu	Ser	Asn	Lys	Asp	Val	Lys	Ile	Glu	Pro	Cys	Gly	His
		355					360					365			
Leu	Leu	Cys	Ser	Cys	Cys	Leu	Ala	Ala	Trp	Gln	His	Ser	Asp	Ser	Gln
	370					375					380				
Thr	Cys	Pro	Phe	Cys	Arg	Cys	Glu	Ile	Lys	Gly	Trp	Glu	Ala	Val	Ser
385				390						395					400
Ile	Tyr	Gln	Phe	His	Gly	Gln	Ala	Thr	Ala	Glu	Asp	Ser	Gly	Asn	Ser
			405					410						415	
Ser	Asp	Gln	Glu	Gly	Arg	Glu	Leu	Glu	Leu	Gly	Gln	Val	Pro	Leu	Ser
		420													

<400> 3

atggctcttg	cggtggc	gtggggcg	cagtgggaag	aggcccg	cctgggccc	60
gcagtcagga	tgctgcagc	cctagaagag	caatgcgtcg	acccccggct	gtccgtgagt	120
cccccttcgc	tgccgggacct	gctgccccgc	acagcgcagc	tgcttcgaga	ggtggcccat	180
tctcggcg	cgcccgccg	aggcgccccc	gggggtccc	gcggctctgg	ggactttcta	240
ctcatctacc	tggccaatct	ggaggccaag	agcaggcagg	tggccgcgct	gctgcctccc	300
cggggccgaa	ggagtgccaa	cgacgagctc	ttccggggcg	gctccagact	caggcgacag	360
ctggccaagc	tggccatcat	cttcagccac	atgcacgcag	agctgcacgc	actcttcccc	420
ggggcaaa	actgtggaca	catgtaccag	ctcaccaagg	ccccgccc	caccttctgg	480
agggaaagt	gcggagccc	gtgtgtgctg	ccctgggctg	agtttgagtc	cctcctgggc	540
acctgccacc	ctgtggaacc	aggctgcaca	gccctggcct	tgccgaccac	cattgacctc	600
acctgcagcg	ggcacgtgtc	catcttcgag	ttcgacgtct	tcaccaggct	ctttcagcca	660
tggccaacac	tcctcaagaa	ctggcagctc	ctggcagtc	accacccagg	ctacatggcc	720
ttcctcacct	atgatgaggt	ccaagagcgt	ctgcaggcct	gcagggacaa	gccaggcagt	780
tacatcttcc	ggcccagctg	tactcgccctg	gggcagtggg	ccatcggtta	tgtgagctca	840
gatggcagca	tcctgcagac	catccctgcc	aacaaacccc	tgtcccagg	gctcctggag	900
ggacagaagg	acggcttcta	cctctaccca	gatggaaaga	cccacaaccc	agacctgact	960
gagctcggcc	aggcagaacc	ccagcagcgc	atccacgtgt	cagaggagca	gctgcagctc	1020
tactgggcca	tggactccac	atttgagctc	tgcaagatct	gtgctgagag	caacaaggat	1080
gtgaagattg	agccgtgcgg	gcacctgctc	tgacgtgct	gcctggctgc	ctggcagcac	1140
tcggacagcc	agacctgccc	cttctgcgc	tgcgagatca	agggctggga	ggccgtgagt	1200
atctaccagt	tcacgggtca	ggctactgct	gaggactcag	ggaacagcag	tgaccaggaa	1260
ggcagggagt	tggagctggg	gcaggtgccc	ctttcggctc	ctccattgcc	cccacggcca	1320
gatctgcccc	ccaggaagcc	cagaaatgcc	cagccgaaag	tgagactcct	aaaggggaac	1380
tcccctccag	ctgcgctggg	accccaggac	cctgcccccg	cc		1422

<210> 4
 <211> 462
 <212> DNA
 <213> Homo Sapiens

tgacagacca	ggcttggcag	tttatttcgg	tttcacaacc	cccttcagc	ccttggggtc	60
ccttgagcag	cacatctggg	tgccctggcc	ttcagcgggn	agnngtcct	ggggtccag	120
cgcangangn	gggagttccc	ctttaggagt	ctcactttcg	gctgggcatt	tctgggcttc	180
ctggggggca	gatctggccg	tgggggcaat	ggaggagccn	aaaggggcac	ctgcccaggc	240
tcacactccc	tgccttccctg	gtcactgctg	ttccctgagt	cctcagcagt	agcctgaccg	300
tagaactgg	agatactcac	ggcctcccag	cccttgatct	cgcagcggca	gaaggggcag	360
gtctgggctg	tcagagtgt	gccaggcanc	caggcagcag	ctgcagaana	ggtgcccgc	420
cggctcaatc	ttcacatcct	tgttgctctc	agcacagatc	tt		462

<210> 5
 <211> 435
 <212> DNA
 <213> Homo Sapiens

agggacaagc	caggcagtta	catcttccgg	cccagctgta	ctcgccctggg	ncagtgggccc	60
atcgntattg	tgagctcaga	tggcagcatc	ctgcagacca	tccctgcca	caaaccctgt	120
tcccagggtg	tcctggagg	acagaaggac	ggcttctacc	tctaccaga	tggaaagacc	180
cacaaccag	acctgactga	gctcggccag	gcagaacccc	agcagcgcat	ccacgtgtca	240
gaggagcagc	tgacgtctta	ctgggcccag	gactccacat	ttgagctctg	caagatctgt	300
gctgagagca	acaaggatgt	gaagattgag	ccgtgcgggc	acctgctctg	cagctgctgc	360
cttggtctgc	cttggcagaa	cttcggacag	ccagaccttg	ccccttcttg	cccgtgcga	420
gattcaagg	cttgg					435

<210> 6

<211> 491
 <212> DNA
 <213> Sus Scrofa

<400> 6

gtgagctcag	atgggcagcat	cctgcagacc	atccctctca	acaaacctct	gttccaagca	60
ctcctggaag	gacaaaagga	aggcttctac	ctctaccctg	atgggaagaa	ccacaacccg	120
gacctgacgg	agctctgcgt	aacgggaccc	tatcaacgca	tccacgtgtc	ggaggagcag	180
ctgcagctgt	actggggccat	gaactccaca	tacgagctct	gcaagatctg	tgccgagaga	240
aacaaggacg	tgaagattga	gccatgcggg	gactgctct	gcagcccgt	gcctgggcta	300
cctggcagaa	ctcagacagn	ccagacctgc	ccttttctgg	ccgctgccag	attcaagggg	360
ccagagcctg	tgngttntcc	atcagtttcc	anggganggc	cagaggggaag	ccgggcgctn	420
aangacncca	nggggacagc	agtggccaag	ggaagatggg	ggatgaggag	ctgggccagg	480
tggacccct c						491

<210> 7
 <211> 2230
 <212> DNA
 <213> C. Elegans

<400> 7

ctatgatcat	tacatcctaa	ttaattgcc	ctggacttca	catcatatca	ccgtttcacc	60
gggaatgggt	tcaataaaca	caatttttca	ccgatacat	cggtttgtca	atggcacagg	120
caataatgcg	cgatttggtc	ccagcacaaa	caactcgacg	gaagcgttga	cactcagtcc	180
gagagctggt	cccagcacag	tttcactatt	cgaaatccca	tcagcttcgg	agatgcccgg	240
tttctgcagt	gaagaggatc	gtcgattttt	gctcaaagca	tgcaagttta	tggatcaagt	300
agtgaagagt	tgatcatagc	caagactgaa	tttgaaaaat	tcgccgcctt	tcattttgga	360
cattctacct	gatacttata	cgcattttaat	gctgatattc	acacaaaaca	atgacatact	420
ccaagacaac	gactacttga	aaatctttct	ggagagtatg	atcaacaagt	gcaaagagat	480
catcaaactg	ttcaagacgt	cagctatcta	caatgaccag	tctgaagaac	gacggaagct	540
tacgaaaatg	tcactaacat	tttcacatat	gcttttcgag	attaaagcat	tatttccgga	600
aggatatctat	attgaagacc	ggtttcggat	gacaaagaag	gaagccgaaa	gcttttggag	660
tcattcatctt	acaaaaaaaa	acattgtacc	ctggtcaaca	ttttttactg	cattagaaaa	720
gcaccatgga	tcaacgatag	gaaaaatgga	agcagccgaa	ttaaaagcta	cgatagactt	780
gagcggagat	gatttttattt	cgaattttga	gtttgatgtg	tttacaaggt	tattctaccc	840
tttcaaaaaca	ctgatcaaaa	attggcaaac	actcaccacc	gcccattccc	gatactgtgc	900
attttctcaca	tacgatgagg	tcaaaaaacg	gttagaaaaa	ttaacgaaaa	aacctggaag	960
ctacatcttc	cggttatcat	gcacacgtcc	tggaacaatg	gcaataggat	acgtagctcc	1020
ggatggaaaag	atttatcaga	caataaccaca	gaataaaaag	ttgattcaag	cactacatga	1080
aggccataaaa	gaaggatttt	atattttacc	gaacggtaga	gatcaagata	ttaacttatc	1140
caaattgatg	gatgtgccac	aagcggacag	agtgcgaagt	accagtgaac	aatacgagtt	1200
gtattgtgag	atgggcacaa	cattcgagtt	gtgcaaaatt	tgtgacgata	acgagaagaa	1260
catcaaaatt	gagccatgtg	gacattttgt	ctgcgcaaaa	tgtttggtta	actggcagga	1320
ttcggatggg	ggtggcaaca	catgtccatt	ctgccgctac	gaaatcaaa	gaacaaatcg	1380
tgtgattatt	gacaggttca	agcccactcc	ggtagaaatt	gaaaaagcga	aaaatgtagc	1440
tgctgcggag	aagaagctga	tctcattagt	tcccgcagtg	cctcccagaa	cgtatgtgtc	1500
ccaatgttct	caaagtttgc	tgcatgacgc	gtcaaaactca	attccgtcgg	tcgacgagtt	1560
gccgttggtg	ccgccaccgt	tgccaccgaa	agcattgggt	accctggaca	ctttgaattc	1620
gtcacaaca	tcctcttcat	acgtgaacat	caaagagctg	gaaaatgttg	aaacaagcgg	1680
agaagcattg	gcacaagtgg	taaaccggca	acgggcgcct	tcaatccaag	ctccaccact	1740
accgccaaag	ttatcagcga	gcgagcacca	accacaccac	ccatacacaa	atacgaacag	1800
tgagcgggag	tagacttggt	taaatgttca	tcttaccgct	ttatactgca	attttcattc	1860
ccccacttat	catagaacta	ttcttccaca	acaacatatt	gccgtgacta	gaactggtaa	1920
cactacatca	ttctttgtta	aaacgttatt	atatctctat	ttctttttcg	cctactcctt	1980
tccgtttttt	tttcaaattt	tgtcaatttt	cctacagcgt	tctgactcct	attggtaagc	2040
aatcatgtca	tatcttggtta	aattttcatg	ttaatttctt	actctcgtg	tcccagattt	2100

tacggagttt	tcaggaaaacg	tttgattttg	ttctattcta	caatttccat	cgcccccaac	2160
ctgtcgtgta	ttttctatgt	gtcactctga	agaaaacaag	tttagacttt	ttaaaaaaaa	2220
aaaaaaaaaa						2230

<210> 8
 <211> 3090
 <212> DNA
 <213> Homo Sapiens

<400> 8						
gaattccggg	cccgatagc	cggcgggcggc	ggcgggcggcg	gcggcgggcgg	cgcccggggag	60
agggccctcc	ttcacgccc	gcttctctcc	ctcgctcgca	gtcgagccga	gccggcgggac	120
ccgcctgggc	tccgaccctg	cccaggccat	ggccggcaac	gtgaagaaga	gctctggggc	180
cgggggcggc	acgggctccg	ggggctcggg	ttcgggtggc	ctgattgggc	tcatgaagga	240
cgccttcag	ccgaccacc	accaccacca	ccacctcagc	ccccacccgc	cggggacggt	300
ggacaagaag	atgggtggaga	agtgtctggaa	gctcatggac	aagggtggtgc	ggttgtgtca	360
gaacccaaag	ctggcgctaa	agaatagccc	accttatatc	ttagacctgc	taccagatac	420
ctaccagcat	ctccgtacta	tcttgtcaag	atatgagggg	aagatggaga	cacttgagga	480
aaatgagtat	tttaggggtg	ttatggagaa	tttgatgaag	aaaactaagc	aaaccataag	540
cctcttcaag	gagggaaaag	aaagaatgta	tgaggagaat	tctcagccta	ggcgaaacct	600
aaccaaactg	tccctcatct	tcagccacat	gctggcagaa	ctaaaaggaa	tctttccaag	660
tggactcttt	cagggagaca	catttcggat	tactaaagca	gatgctgcgg	aattttggag	720
aaaagctttt	ggggaaaaga	caatagtccc	ttggaagagc	tttcgacagg	ctctacatga	780
agtgcacccc	atcagttctg	ggctggaggc	catggctctg	aaatccacta	ttgatctgac	840
ctgcaatgat	tatatctcgg	tttttgaaat	tgacatcttt	acccgactct	ttcagccctg	900
gtcctctttg	ctcaggaatt	ggaacagcct	tgctgtaact	catcctggct	acatggcttt	960
tttgacgtat	gacgaagtga	aagctcggct	ccagaaattc	attcacaaac	ctggcagtta	1020
tatcttccgg	ctgagctgta	ctcgtctggg	tcagtgggct	attgggtatg	ttactgctga	1080
tgggaacatt	ctccagacaa	tccctcacaa	taaacctctc	ttccaagcac	tgattgatgg	1140
cttcagggaa	ggcttctatt	tgtttcctga	tggacgaaat	cagaatcctg	atctgactgg	1200
cttatgtgaa	ccaactcccc	aagaccatat	caaagtgacc	caggaacaat	atgaattata	1260
ctgtgagatg	ggctccacat	tccaactatg	taaaatatgt	gctgaaaatg	ataaggatgt	1320
aaagattgag	ccctgtggac	acctcatgtg	cacatcctgt	cttacatcct	ggcaggaatc	1380
agaaggctcag	ggctgtcctt	tctgccgatg	tgaaattaaa	ggtactgaac	ccatcgtggt	1440
agatccgttt	gatccatagag	ggagtggcag	cctgttgagg	caaggagcag	agggagctcc	1500
ctcccaaat	tatgatgatg	atgatgatga	acgagctgat	gatactctct	tcatgatgaa	1560
ggaattggct	ggtgccaaag	tggaaaggcc	gccttctcca	ttctccatgg	ccccacaagc	1620
ttcccttccc	ccggtgccac	cacgacttga	ccttctgccg	cagcgagtat	gtgttccctc	1680
aagtgttct	gctcttgga	ctgcttctaa	ggctgttct	ggctcccttc	ataaagacaa	1740
accattgcca	gtacctcca	cacttcgaga	tcttccacca	ccaccgcctc	cagaccggcc	1800
atattctgtt	ggagcagaat	cccgaacctca	aagacgcccc	ttgccttgta	caccaggcga	1860
ctgtccctcc	agagacaaac	tgccccctgt	ccctcttagc	cgccttggag	actcatggct	1920
gccccggcca	atccccaag	taccagtatc	tgccccaagt	tccagtgatc	cctggacagg	1980
aagagaatta	accaaccggc	actcacttcc	atthttcattg	ccctcacaaa	tggagcccag	2040
accagatgtg	cctaggctcg	gaagcacgtt	cagtctggat	acctccatga	gtatgaatag	2100
cagccatta	gtaggctcag	agtgtgacca	ccccaaaatc	aaaccttct	catctgccaa	2160
tgccatttat	tctctggctg	ccagacctct	tcctgtgcca	aaactgccac	ctggggagca	2220
atgtgagggt	gaagaggaca	cagagtacat	gactccctct	tccaggcctc	tacggccttt	2280
ggatacatcc	cagagttcac	gagcatgtga	ttgcgaccag	cagattgata	gctgtacgta	2340
tgaagcaatg	tataatattc	agtcccaggc	gccatctatc	accgagagca	gcaccttttg	2400
tgaagggaat	ttggccgcag	cccatgccaa	cactgggtccc	gaggagtcag	aaaatgagga	2460
tgatgggtat	gatgtcccaa	agccacctgt	gccggccgtg	ctggcccgcc	gaactctctc	2520
agatatctct	aatgccagct	cctccttttg	ctggttgtct	ctggatgggtg	atcctacaac	2580
aaatgtcact	gaaggttccc	aagttcccga	gaggcctcca	aaaccattcc	cgcggagaat	2640
caactctgaa	cggaaagctg	gcagctgtca	gcaaggtagt	ggtcctgcgc	cctctgctgc	2700
caccgcctca	cctcagctct	ccagtgaagt	cgagaacctc	atgagtcagg	ggtactccta	2760

ccaggacatc	cagaaagctt	tgggtcattgc	ccagaacaac	atcgagatgg	ccaaaaacat	2820
cctccgggaa	tttgtttcca	tttcttctcc	tgcccatgta	gctacctagc	acaccatctc	2880
cctgctgcag	gttttagagga	ccagtgagtt	gggagttatt	actcaagtgg	cacctagaag	2940
ggcaggagtt	ccttttggtga	cttcacagtg	aagtcttgcc	ctctctgtgg	gatatcacat	3000
cagtggttcc	aagatttcaa	agtggtgaaa	tgaaaatgga	gcagctagta	tgttttatta	3060
ttttatgggt	cttgagtgca	tttgaagggt				3090

<210> 9
 <211> 3982
 <212> DNA
 <213> Homo Sapiens

<400> 9						
ctgggtcctg	tgtgtgccac	aggggtgggg	tgtccagcga	gcggtctect	cctcctgeta	60
gtgctgctgc	ggcgtcccgc	ggcctccccg	agtcggggcg	gaggggagag	cggtgtgga	120
tttgtcttga	cggttaattgt	tgcgtttcca	cgtctcggag	gcctgcgcgc	tgggttgctc	180
cttcttcggg	agcagctgtg	tctcagcgat	cccactcca	gccggggctc	cccacacaca	240
ctgggctgga	tgcgtgtgga	gtgggacccg	cgcacacgcg	tgtctctgga	cagctacggc	300
gccgaaagaa	ctaaaattcc	agatggcaaa	ctcaatgaat	ggcagaaacc	ctgggtggtcg	360
aggaggaaat	ccccgaaaag	gtcgaatttt	gggtattatt	gatgctattc	aggatgcagt	420
tggacccccct	aagcaagctg	cgcagatcg	caggaccgtg	gagaagactt	ggaagctcat	480
ggacaaagtg	gtaagactgt	gccaaaatcc	caaacttcag	ttgaaaaata	gccaccata	540
tatacttgat	atthttgctg	atacatatca	gcatttacga	cttatattga	gtaaatatga	600
tgacaaccag	aaacttgccc	aactcagtga	gaatgagtag	tttaaaatct	acattgatag	660
ccttatgaaa	aagtcaaaac	gggcaataag	actctttaaa	gaaggcaagg	agagaatgta	720
tgaagaacag	tcacaggaca	gacgaaatct	cacaaaactg	tcccttatct	tcagtcacat	780
gctggcagaa	atcaaagcaa	tctttcccaa	tgggtcaattc	cagggagata	actttcgtat	840
cacaaaagca	gatgctgctg	aattctggag	aaagtttttt	ggagacaaaa	ctatcgtacc	900
atggaaagta	ttcagacagt	gccttcctga	ggccaccag	attagctcta	gcctggaagc	960
aatggctcta	aaatcaacaa	ttgatthaac	ttgcaatgat	tacatttcag	tttttgaatt	1020
tgatattttt	accaggctgt	ttcagccttg	gggctctatt	ttgcggaatt	ggaatttctt	1080
agctgtgaca	catccagggt	acatggcatt	tctcacatat	gatgaagtta	aagcacgact	1140
acagaaatat	agcaccaaaac	cgggaagcta	tattttccgg	ttaagttgca	ctcgattggg	1200
acagtgggoc	attggctatg	tgactgggga	tggaatatc	ttacagacca	tacctcataa	1260
caagccetta	tttcaagccc	tgattgatgg	cagcagggaa	ggattttatc	tttatcctga	1320
tgggaggagt	tataatcctg	atttaactgg	attatgtgaa	cctacacctc	atgaccatat	1380
aaaagttaca	caggaacaat	atgaattata	ttgtgaaatg	ggctccactt	ttcagctctg	1440
taagatttgt	gcagagaatg	acaaagatgt	caagattgag	ccttgtgggc	atttgatgtg	1500
cacctcttgc	cttacggcat	ggcaggagtc	ggatggctag	ggctgccctt	tctgtcgttg	1560
tgaaataaaa	ggaactgagc	ccataatcgt	ggaccccttt	gatccaagag	atgaaggctc	1620
cagggtgttg	agcatcattg	accccttttg	catgccgatg	ctagacttgg	acgacgatga	1680
tgatcgtgag	gagtccttga	tgatgaatcg	gttggcaaac	gtccgaaagt	gcactgacag	1740
gcagaactca	ccagtcacat	caccaggatc	ctctcccctt	gccagagaa	gaaagccaca	1800
gcctgaccca	ctccagatcc	cacatctaag	cctgccaccc	gtgcctcctc	gcctggatct	1860
aattcagaaa	ggcatagtta	gatctcccctg	tggcagccca	acagggttcac	caaagtcttc	1920
tccttgcatg	gtgagaaaaac	aagataaaac	actcccagca	ccacctcctc	ccttaagaga	1980
tcctcctcca	ccgccacctg	aaagacctcc	accaatccca	ccagacaata	gactgagtag	2040
acacatccat	catgtggaaa	gcgtgccttc	cagagaccgg	ccaatgcctc	ttgaagcatg	2100
gtgcccctgg	gatgtgtttg	ggactaatca	gcttgtggga	tgctgactcc	taggggaggg	2160
ctctccaaaa	cctggaatca	cagcgagttc	aaatgtccaat	ggaaggcaca	gtagagtggg	2220
ctctgaccca	gtgcttatgc	ggaaacacag	acgccatgat	ttgccttttag	aaggagctaa	2280
gggtcttttcc	aatgggtcac	ttggaagtga	agaatatgat	gttcctcccc	ggctttctcc	2340
tcctcctcca	gttaccaccc	tcctccctag	cataaagtgt	actgggtccgt	tagcaaattc	2400
tccttcagag	aaaacaagag	acccagtaga	ggaagatgat	gatgaatata	agattccttc	2460
atccacccct	gtttccctga	attcacaacc	atctcattgt	cataatgtaa	aacctcctgt	2520
tcgggtcctgt	gataatgggc	actgtatgct	gaatggaaca	catgggtccat	cttcagagaa	2580

gaaatcaaac	atccctgact	taagcatata	tttaaagggga	gatgtttttg	attcagcctc	2640
tgatcccgtg	ccattaccac	ctgccaggcc	tccaactcgg	gacaatccaa	agcatgggtc	2700
ttcactcaac	aggacgccct	ctgattatga	tcttctcatc	cctccattag	gtgaagatgc	2760
ttttgatgcc	ctccctccat	ctctcccacc	tccccacct	cctgcaaggc	atagtctcat	2820
tgaacattca	aaacctcctg	gctccagtag	ccggccatcc	tcaggacagg	atctttttct	2880
tcttccttca	gatccctttg	ttgatctagc	aagtggccaa	gttcctttgc	ctcctgctag	2940
aagggtacca	ggtgaaaatg	tcaaaactaa	cagaacatca	caggactatg	atcagcttcc	3000
ttcatgttca	gatgggttcac	aggcaccagc	cagacccctt	aaaccacgac	cgcgaggagc	3060
tgcaccagaa	attcaccaca	gaaaacccca	tgggcctgag	gcggcattgg	aaaatgtcga	3120
tgcaaaaatt	gcaaaactca	tgggagaggg	ttatgccttt	gaagaggtga	agagagcctt	3180
agagatagcc	cagaataatg	tcgaagttgc	ccggagcatc	ctccgagaat	ttgccttccc	3240
tctccagta	tccccacgtc	taaatctata	gcagccagaa	ctgtagacac	caaaatggaa	3300
agcaatcgat	gtattccaag	agtgtggaaa	taaagagaac	tgagatggaa	ttcaagagag	3360
aagtgtctcc	tctcgtgtga	gcagcttgag	aagaggcttg	ggagtgcagc	ttctcaaagg	3420
agacogatgc	ttgctcagga	tgtcgacagc	tgtggcttcc	ttgtttttgc	tagccatatt	3480
tttaaatacag	ggttgaactg	acaaaaataa	tttaaagacg	tttacttccc	ttgaactttg	3540
aacctgtgaa	atgctttacc	ttgttttaca	tttggcaaag	ttgcagtttg	ttcttgtttt	3600
tagtttagtt	ttgttttggg	gttttgatac	ctgtactgtg	ttcttcacag	accctttgta	3660
gcgtgggtcag	gtctgctgta	acatttccca	ccaactctct	tgctgtccac	atcaacagct	3720
aaatcattta	ttcatatgga	tctctaccat	ccccatgcct	tgcccaggtc	cagttccatt	3780
tctctcattc	acaagatgct	ttgaagggtc	tgattttcaa	ctgatcaaac	taatgcaaaa	3840
aaaaaaaagta	tgtattcttc	actactgagt	ttcttctttg	gaaaccatca	ctattgagag	3900
atgggaaaaa	cctgaatgta	taaagcattt	atttgtcaat	aaactgcctt	ttgtaagggg	3960
ttttcacata	aaaaaaaaaa	aa				3982

<210> 10
 <211> 17
 <212> PRT
 <213> Homo Sapiens

<400> 10
 Arg Lys Pro Arg Asn Ala Gln Pro Lys Val Arg Leu Leu Lys Gly Asn
 1 5 10 15
 Ser

SEQUENCE LISTING

<110> Borriello, Francescopaolo
 Band, Hamid

<120> Characterization of a novel gene Cbl-SL

<130> B0801/7159/ERP

<150> U.S. 60/107,470
 <151> 1998-11-06

<160> 10

<170> FastSEQ for Windows Version 3.0

<210> 1
 <211> 1547
 <212> DNA
 <213> Homo Sapiens

<220>
 <221> CDS
 <222> (13)...(1434)

<400> 1
 cgcgaggctc cc atg gct ctg gcg gtg gcc ccg tgg ggg cga cag tgg gaa 51
 Met Ala Leu Ala Val Ala Pro Trp Gly Arg Gln Trp Glu
 1 5 10

gag gcc cgc gcc ctg ggc cgg gca gtc agg atg ctg cag cgc cta gaa 99
 Glu Ala Arg Ala Leu Gly Arg Ala Val Arg Met Leu Gln Arg Leu Glu
 15 20 25

gag caa tgc gtc gac ccc cgg ctg tcc gtg agt ccc cct tgc ctg cgg 147
 Glu Gln Cys Val Asp Pro Arg Leu Ser Val Ser Pro Pro Ser Leu Arg
 30 35 40 45

gac ctg ctg ccc cgc aca gcg cag ctg ctt cga gag gtg gcc cat tct 195
 Asp Leu Leu Pro Arg Thr Ala Gln Leu Leu Arg Glu Val Ala His Ser
 50 55 60

cgg cgg gcg gcc ggc gga ggc ggc ccc ggg ggt ccc ggc ggc tct ggg 243
 Arg Arg Ala Ala Gly Gly Gly Gly Pro Gly Gly Pro Gly Gly Ser Gly
 65 70 75

gac ttt cta ctc atc tac ctg gcc aat ctg gag gcc aag agc agg cag 291
 Asp Phe Leu Leu Ile Tyr Leu Ala Asn Leu Glu Ala Lys Ser Arg Gln
 80 85 90

gtg gcc gcg ctg ctg cct ccc cgg ggc cga agg agt gcc aac gac gag 339
 Val Ala Ala Leu Leu Pro Pro Arg Gly Arg Arg Ser Ala Asn Asp Glu
 95 100 105

655011 B0246460

ctc ttc cgg gcg ggc tcc aga ctc agg cga cag ctg gcc aag ctg gcc Leu Phe Arg Ala Gly Ser Arg Leu Arg Arg Gln Leu Ala Lys Leu Ala 110 115 120 125	387
atc atc ttc agc cac atg cac gca gag ctg cac gca ctc ttc ccc ggg Ile Ile Phe Ser His Met His Ala Glu Leu His Ala Leu Phe Pro Gly 130 135 140	435
gca aag tac tgt gga cac atg tac cag ctc acc aag gcc ccc gcc cac Ala Lys Tyr Cys Gly His Met Tyr Gln Leu Thr Lys Ala Pro Ala His 145 150 155	483
acc ttc tgg agg gaa agt tgc gga gcc cgg tgt gtg ctg ccc tgg gct Thr Phe Trp Arg Glu Ser Cys Gly Ala Arg Cys Val Leu Pro Trp Ala 160 165 170	531
gag ttt gag tcc ctc ctg ggc acc tgc cac cct gtg gaa cca ggc tgc Glu Phe Glu Ser Leu Leu Thr Cys His Pro Val Glu Pro Gly Cys 175 180 185	579
aca gcc ctg gcc ttg cgc acc acc att gac ctc acc tgc agc ggg cac Thr Ala Leu Ala Leu Arg Thr Thr Ile Asp Leu Thr Cys Ser Gly His 190 195 200 205	627
gtg tcc atc ttc gag ttc gac gtc ttc acc agg ctc ttt cag cca tgg Val Ser Ile Phe Glu Phe Asp Val Phe Thr Arg Leu Phe Gln Pro Trp 210 215 220	675
cca aca ctc ctc aag aac tgg cag ctc ctg gca gtc aac cac cca ggc Pro Thr Leu Leu Lys Asn Trp Gln Leu Leu Ala Val Asn His Pro Gly 225 230 235	723
tac atg gcc ttc ctc acc tat gat gag gtc caa gag cgt ctg cag gcc Tyr Met Ala Phe Leu Thr Tyr Asp Glu Val Gln Glu Arg Leu Gln Ala 240 245 250	771
tgc agg gac aag cca ggc agt tac atc ttc cgg ccc agc tgt act cgc Cys Arg Asp Lys Pro Gly Ser Tyr Ile Phe Arg Pro Ser Cys Thr Arg 255 260 265	819
ctg ggg cag tgg gcc atc ggc tat gtg agc tca gat ggc agc atc ctg Leu Gly Gln Trp Ala Ile Gly Tyr Val Ser Ser Asp Gly Ser Ile Leu 270 275 280 285	867
cag acc atc cct gcc aac aaa ccc ctg tcc cag gtg ctc ctg gag gga Gln Thr Ile Pro Ala Asn Lys Pro Leu Ser Gln Val Leu Leu Glu Gly 290 295 300	915
cag aag gac ggc ttc tac ctc tac cca gat gga aag acc cac aac cca Gln Lys Asp Gly Phe Tyr Leu Tyr Pro Asp Gly Lys Thr His Asn Pro 305 310 315	963
gac ctg act gag ctc ggc cag gca gaa ccc cag cag cgc atc cac gtg Asp Leu Thr Glu Leu Gly Gln Ala Glu Pro Gln Gln Arg Ile His Val 320 325 330	1011

tca gag gag cag ctg cag ctc tac tgg gcc atg gac tcc aca ttt gag	1059
Ser Glu Glu Gln Leu Gln Leu Tyr Trp Ala Met Asp Ser Thr Phe Glu	
335 340 345	
ctc tgc aag atc tgt gct gag agc aac aag gat gtg aag att gag ccg	1107
Leu Cys Lys Ile Cys Ala Glu Ser Asn Lys Asp Val Lys Ile Glu Pro	
350 355 360 365	
tgc ggg cac ctg ctc tgc agc tgc tgc ctg gct gcc tgg cag cac tcg	1155
Cys Gly His Leu Leu Cys Ser Cys Cys Leu Ala Ala Trp Gln His Ser	
370 375 380	
gac agc cag acc tgc ccc ttc tgc cgc tgc gag atc aag ggc tgg gag	1203
Asp Ser Gln Thr Cys Pro Phe Cys Arg Cys Glu Ile Lys Gly Trp Glu	
385 390 395	
gcc gtg agt atc tac cag ttc cac ggt cag gct act gct gag gac tca	1251
Ala Val Ser Ile Tyr Gln Phe His Gly Gln Ala Thr Ala Glu Asp Ser	
400 405 410	
ggg aac agc agt gac cag gaa ggc agg gag ttg gag ctg ggg cag gtg	1299
Gly Asn Ser Ser Asp Gln Glu Gly Arg Glu Leu Glu Leu Gly Gln Val	
415 420 425	
ccc ctt tcg gct cct cca ttg ccc cca cgg cca gat ctg ccc ccc agg	1347
Pro Leu Ser Ala Pro Pro Leu Pro Pro Arg Pro Asp Leu Pro Pro Arg	
430 435 440 445	
aag ccc aga aat gcc cag ccg aaa gtg aga ctc cta aag ggg aac tcc	1395
Lys Pro Arg Asn Ala Gln Pro Lys Val Arg Leu Leu Lys Gly Asn Ser	
450 455 460	
cct cca gct gcg ctg gga ccc cag gac cct gcc ccg gcc tgaaggccag	1444
Pro Pro Ala Ala Leu Gly Pro Gln Asp Pro Ala Pro Ala	
465 470	
ggcaccacaga tgtgtgtgtc aagggagccc caagggctgg aaggggggtg tgaaaccgaa	1504
ataaactgcc aagcctgggtc tgtcaaaaaa aaaaaaaaaa aaa	1547

<210> 2

<211> 474

<212> PRT

<213> Homo Sapiens

<400> 2

Met Ala Leu Ala Val Ala Pro Trp Gly Arg Gln Trp Glu Glu Ala Arg	
1 5 10 15	
Ala Leu Gly Arg Ala Val Arg Met Leu Gln Arg Leu Glu Glu Gln Cys	
20 25 30	
Val Asp Pro Arg Leu Ser Val Ser Pro Pro Ser Leu Arg Asp Leu Leu	
35 40 45	
Pro Arg Thr Ala Gln Leu Leu Arg Glu Val Ala His Ser Arg Arg Ala	
50 55 60	
Ala Gly Gly Gly Gly Pro Gly Gly Pro Gly Gly Ser Gly Asp Phe Leu	
65 70 75 80	
Leu Ile Tyr Leu Ala Asn Leu Glu Ala Lys Ser Arg Gln Val Ala Ala	

atggctctgg	cggtggcccc	gtggggggcga	cagtgggaag	aggccccgcgc	cctggggccgg	60
gcagtcagga	tgctgcagcg	cctagaagag	caatgcgtcg	acccccggct	gtccgtgagt	120
cccccttcgc	tgcgggacct	gctgccccgc	acagcgcagc	tgcttcgaga	ggtggcccat	180
tctcggcggg	cggccggcgg	aggcgcccc	gggggtcccg	gcggctctgg	ggactttcta	240
ctcatctacc	tggccaatct	ggaggccaag	agcaggcagg	tggccgcgct	gctgcctccc	300
cggggccgaa	ggagtgccaa	cgacgagctc	ttccggggcg	gctccagact	caggcgacag	360
ctggccaagc	tggccatcat	cttcagccac	atgcacgcag	agctgcacgc	actcttcccc	420
ggggcaaagt	actgtggaca	catgtaccag	ctcaccaagg	ccccgccca	caccttctgg	480
agggaaagt	gcggagcccg	gtgtgtgctg	ccctgggctg	agtttgagtc	cctcctgggc	540
acctgccacc	ctgtggaacc	aggctgcaca	gccctggcct	tgcgaccac	cattgacctc	600
acctgcagcg	ggcacgtgtc	catcttcgag	ttcgacgtct	tcaccaggct	ctttcagcca	660
tggccaacac	tcctcaagaa	ctggcagctc	ctggcagtca	accaccagg	ctacatggcc	720
ttcctcacct	atgatgaggt	ccaagagcgt	ctgcaggcct	gcagggacaa	gccaggcagt	780
tacatcttcc	ggcccagctg	tactcgctcg	gggcagtggg	ccatcggcta	tgtgagctca	840
gatggcagca	tcctgcagac	catccctgcc	aacaaacccc	tgtcccagg	gctcctggag	900
ggacagaagg	acggcttcta	cctctaccca	gatggaaaga	cccacaaccc	agacctgact	960
gagctcggcc	aggcagaacc	ccagcagcgc	atccacgtgt	cagaggagca	gctgcagctc	1020
tactggggcca	tggactccac	atttgagctc	tgcaagatct	gtgctgagag	caacaaggat	1080
gtgaagattg	agccgtgcgg	gcacctgctc	ttcagctgct	gcctggctgc	ctggcagcac	1140
tcggacagct	agacctgccc	cttctgccc	tgcgagatca	agggtggga	ggcgtgagt	1200
atctaccagt	tcacaggctc	ggctactgct	gaggactcag	ggaacagcag	tgaccaggaa	1260
ggcagggagt	tggagctggg	gcaggtgccc	ctttcggtct	ctccattgcc	cccacggcca	1320
gatctgcccc	ccaggaagcc	cagaaatgcc	cagccgaaag	tgagactcct	aaaggggaac	1380
tccctccag	ctgcgctggg	accccaggac	cctgccccgg	cc		1422

<210> 4

<211> 462

<212> DNA

<213> Homo Sapiens

<400> 4

tgacagacca	ggcttggcag	tttatattcgg	tttcacaacc	cccttccagc	ccttgggggtc	60
ccttgagcag	cacatctggg	tgccctggcc	ttcagcgggn	agngngtcct	ggggteccag	120
cgcangan	gggagttccc	ctttaggagt	ctcactttcg	gctgggcatt	tctgggcttc	180
ctggggggca	gatctggccg	tgggggcaat	ggaggagccn	aaaggggcac	ctgcccaggc	240
tccaaactccc	tgcttctcctg	gtcactgctg	ttccctgagt	cctcagcagt	agcctgaccg	300
tagaactggg	agatactcac	ggcctcccag	cccttgatct	cgcagcggca	gaaggggcag	360
gtctgggctg	tcagagtgtc	gccaggcanc	caggcagcag	ctgcagaana	ggtgcccgcga	420
cggctcaatc	ttcacatcct	tgttgctctc	agcacagatc	tt		462

<210> 5

<211> 435

<212> DNA

<213> Homo Sapiens

<400> 5

agggacaagc	caggcagtta	catcttccgg	cccagctgta	ctcgctggg	ncagtggggcc	60
atcgntattg	tgagctcaga	tggcagcatc	ctgcagacca	tccttgccaa	caaacccttg	120
tcccaggtgc	tcctggaggg	acagaaggac	ggcttctacc	tctaccaga	tggaaagacc	180
cacaaccacg	acctgactga	gctcgccag	gcagaacccc	agcagcgcac	ccagctgtca	240
gaggagcagc	tgcagctcta	ctgggccatg	gactccacat	ttgagctctg	caagatctgt	300
gctgagagca	acaaggatgt	gaagattgag	ccgtgcgggc	acctgctctg	cagctgctgc	360
cttggcttgc	cttggcagaa	cttcggacag	ccagaccttg	ccccttcttg	cccgtgcca	420
gattcaaggg	cttgg					435

<210> 6

<211> 491
 <212> DNA
 <213> Sus Scrofa

<400> 6
 gtgagctcag atggcagcat cctgcagacc atccctctca acaaacctct gttccaagca 60
 ctcttggaag gacaaaagga aggtctctac ctctaccctg atgggaagaa ccacaacccg 120
 gacctgacgg agctctgcgt aacgggaccc tatcaacgca tccacgtgtc ggaggagcag 180
 ctgcagctgt actgggccat gaactccaca tacgagctct gcaagatctg tgccgagaga 240
 aacaaggacg tgaagattga gccatgcggg gcactgctct gcagcccgtc gcctgggcta 300
 cctggcagaa ctacagacagn ccagacctgc ccttttctgg ccgctgccag attcaagggg 360
 ccagagcctg tngtntntcc atcagtttcc anggganggc cagaggggaag ccgggcgctn 420
 aangacncca nggggacagc agtggccaag ggaagatggg ggatgaggag ctggggccagg 480
 tggaccccct c 491

<210> 7
 <211> 2230
 <212> DNA
 <213> C. Elegans

<400> 7
 ctatgatcat tacatcctaa ttaattgcc a ctggacttca catcatatca ccgtttcacc 60
 gggaaatggg tcaataaaca caatttttca ccggatacat cggtttgtca atggcacagg 120
 caataatgcg cgatttggtc ccagcacaaa caactcgacg gaagcgttga cactcagtc 180
 gagagctgtt cccagcacag tttcactatt cgaaatccca tcagcttcgg agatgcccg 240
 tttctgcagt gaagaggatc gtcgattttt gctcaaagca tgcaagttta tggatcaagt 300
 agtgaagagt tgtcatagcc caagactgaa tttgaaaaat tcgccgcctt tcattttgga 360
 cattctacct gatacttata cgcattttaat gctgatatc acacaaaaca atgacatact 420
 ccaagacaac gactacttga aaatctttct ggagagtatg atcaacaagt gcaaagagat 480
 catcaaactg ttcaagacgt cagctatcta caatgaccag tctgaagaac gacggaagct 540
 tacgaaaatg tctaatacat tttcacatat gcttttcgag attaaagcat tatttccgga 600
 aggtatctat attgaagacc ggttttcggat gacaaaagaag gaagccgaaa gcttttggag 660
 tcatcatttt acaaaaaaaaa acattgtacc ctggtcaaca ttttttactg cattagaaaa 720
 gcaccatgga tcaacgatag gaaaaatgga agcagccgaa ttaaaaagcta cgatagactt 780
 gagcggagat gattttattt cgaattttga gtttgatgtg tttacaagg tttctatccc 840
 tttcaaaaaca ctgatcaaaa attggcaaac actcaccacc gcccatcccg gatactgtgc 900
 attttctcaca tacgatgagg tcaaaaaacg gttagaaaaa ttaacgaaaa aacctggaag 960
 ctacatcttc cggttatcat gcacacgtcc tggacaatgg gcaataggat acgtagctcc 1020
 ggatggaaaag atttatcaga caataccaca gaataaaaagt ttgattcaag cactacatga 1080
 aggccataaa gaaggatttt atattttacc gaacggtaga gatcaagata ttaacttatt 1140
 caaattgatg gatgtgccac aagcggacag agtgcaagt accagtgaac aatacgagtt 1200
 gtattgtgag atgggcacaa cattcgagtt gtgcaaaatt tgtgacgata acgagaagaa 1260
 catcaaaatt gagccatgtg gacatttgct ctgcgcaaaa tgtttggcta actggcagga 1320
 ttcggatggg ggtggcaaca catgtccatt ctgccgtac gaaatcaaag gaacaaatcg 1380
 tgtgattatt gacaggttca agcccactcc ggtagaaatt gaaaaagcga aaaatgtagc 1440
 tgctgcgag aagaagctga tctcattagt tcccagctg cctcccagaa cgtatgtgtc 1500
 ccaatgttct caaagtttgc tgcagacgc gtcaaactca attcgcgtcg tcgacgagtt 1560
 gccgttgggt ccgccaccgt tggcaccgaa agcattgggt accctggaca ctttgaattc 1620
 gtcacaaaaca tctcttcat acgtgaacat caaagagctg gaaaatgttg aaacaagcgg 1680
 agaagcattg gcacaagtgg taaaccggca acgggcgcct tcaatccaag ctccaccact 1740
 accgccaagg ttatcagcga gcgagcacca accacccac ccatcacaa atacgaacag 1800
 tgagcgggag tagacttggt taaatgttca tcttaccgct ttatactgca attttcttc 1860
 cccacttat catagaacta ttcttccaca acaacatatt gccgtgacta gaactggtaa 1920
 cactacatca ttctttgtta aaacgttatt atatctctat ttctttttcg cctactcctt 1980
 tccgtttttt tttcaaat tgtcaatttt cctacagcgt tctgactcct attggtaagc 2040
 aatcatgtca tatcttggtta aattttcatg ttaatttctt actctcgctg tcccagattt 2100

tacggagttt	tcaggaaacg	tttgattttg	ttctattcta	caattttccat	cgcccccaac	2160
ctgtcgtgta	ttttctatgt	gtcactctga	agaaaacaag	tttagacttt	ttaaaaaaaa	2220
aaaaaaaaaa						2230

<210> 8
 <211> 3090
 <212> DNA
 <213> Homo Sapiens

<400> 8

gaattccggg	cccgatagc	cggcggcggc	ggcgggcgcg	gcgggcgcg	cgcccgggag	60
aggccctcc	ttcacgcct	gcttctctcc	ctcgtctcgca	gtcgagccga	gccggcgggac	120
ccgctgggc	tccgacctg	cccaggccat	ggccggcaac	gtgaagaaga	gctctggggc	180
cggggcggc	acgggtccg	ggggtcggg	ttcggtggc	ctgattgggc	tcatgaagga	240
cgcttccag	ccgcaccacc	accaccacca	ccacctcagc	ccccaccgc	cggggacggg	300
ggacaagaag	atggtggaga	agtgtggaa	gctcatggac	aagggtggtgc	ggttgtgtca	360
gaacccaaag	ctggcgctaa	agaatagccc	accttatatc	ttagacctgc	taccagatac	420
ctaccagcat	ctccgtacta	ctttgtcaag	atatgagggg	aagatggaga	cacttggaga	480
aaatgagtat	tttaggggtg	ttatggagaa	ttgatgaag	aaaactaagc	aaaccataag	540
cctcttcaag	gagggaag	aaagaatgta	tgaggagaat	tctcagccta	ggcgaaacct	600
aaccaaactg	tccctcatct	tcagccacat	gctggcagaa	ctaaaaggaa	tctttccaag	660
tggactcttt	caggagaca	catttcggat	tactaaagca	gatgctgcgg	aattttggag	720
aaaagctttt	ggggaaaaga	caatagtccc	ttggaagagc	tttcgacagg	ctctacatga	780
agtgcacccc	atcagttctg	ggctggaggc	catggctctg	aaatccacta	ttgatctgac	840
ctgcaatgat	tatatttcgg	tttttgaatt	tgacatcttt	acccgactct	ttcagccctg	900
gtcctctttg	ctcaggaatt	ggaacagcct	tgctgtaact	catcctggct	acatggcttt	960
tttgacgtat	gacgaagtga	aagctcggct	ccagaaattc	attcacaac	ctggcagtta	1020
tatcttccgg	ctgagctgta	ctcgtctggg	tcagtgggct	attgggtatg	ttactgctga	1080
tgggaacatt	ctccagacaa	tccctcacia	taaacctctc	ttccaagcac	tgattgatgg	1140
cttcagggaa	ggcttctatt	tgtttcctga	tggacgaaat	cagaatcctg	atctgactgg	1200
cttatgtgaa	ccaactcccc	aagaccatat	caaagtgacc	caggaacaat	atgaattata	1260
ctgtgagatg	ggctccacat	tccaactatg	taaaatatgt	gctgaaaatg	ataaggatgt	1320
aaagattgag	ccctgtggac	acctcatgtg	cacatcctgt	cttacatcct	ggcaggaatc	1380
agaaggtcag	ggctgtcctt	tctgccgatg	tgaaattaaa	ggtaactgaac	ccatcgtggt	1440
agatccgttt	gatcctagag	ggagtggcag	cctgttgagg	caaggagcag	agggagctcc	1500
ctccccaaat	tatgatgatg	atgatgatga	acgagctgat	gatactctct	tcgatgatga	1560
ggaattggct	ggtgccaaag	tggaaacggc	gccttctcca	ttctccatgg	ccccacaagc	1620
ttcccttccc	ccggtgccac	cacgacttga	ccttctgcgg	cagcgagtat	gtgttccctc	1680
aagtgtctct	gctcttggaa	ctgcttctaa	ggctgtctct	ggctcccttc	ataaagacaa	1740
accattgcca	gtacctccca	cacttcgaga	tcttccacca	ccaccgcctc	cagaccggcc	1800
atattctgtt	ggagcagaat	cccgacctca	aagacgcccc	ttgccttgta	caccaggcga	1860
ctgtccctcc	agagacaaac	tgccccctgt	cccccttagc	cgccttggag	actcatggct	1920
gccccggcca	atccccaaag	taccagtatc	tgccccaaag	tccagtgatc	cctggacagg	1980
aagagaatta	accaaccggc	actcacttcc	atthttcattg	ccctcacaaa	tggagcccag	2040
accagatgtg	cctaggctcg	gaagcacgtt	cagtctggat	acctccatga	gtatgaatag	2100
cagcccatga	gtagggtccag	agtgtgacca	ccccaaaatc	aaaccttcc	catctgccaa	2160
tgccatttat	tctctggctg	ccagacctct	tcctgtgcca	aaactgccac	ctggggagca	2220
atgtgagggg	gaagaggaca	cagagtacat	gactccctct	tccaggcctc	tacggccttt	2280
ggatacatcc	cagagttcac	gagcatgtga	ttgcgaccag	cagattgata	gctgtacgta	2340
tgaagcaatg	tataatatcc	agtcaccagg	gccatctatc	accgagagca	gcacctttgg	2400
tgaagggaaat	ttggcccgag	cccatgccaa	cactggtccc	gaggagtcag	aaaatgagga	2460
tgatgggtat	gatgtcccaa	agccacctgt	gccggccgtg	ctggcccggc	gaactctctc	2520
agatatctct	aatgccagct	cctcctttgg	ctggttgtct	ctggatgggtg	atcctacaac	2580
aaatgtcact	gaaggttccc	aagttcccca	gaggcctcca	aaaccattcc	cgcggagaat	2640
caactctgaa	cggaaagctg	gcagctgtca	gcaaggtagt	ggctcctgcc	cctctgctgc	2700
caccgcctca	cctcagctct	ccagtgaagt	cgagaacctc	atgagtcagg	ggtactccta	2760

ccaggacatc	cagaaagctt	tgggtcattgc	ccagaacaac	atcgagatgg	ccaaaaacat	2820
cctccgggaa	tttgtttcca	tttcttctcc	tgcccatgta	gctacctagc	acaccatctc	2880
cctgctgcag	gttttagagga	ccagtgagtt	gggagttatt	actcaagtgg	cacctagaag	2940
ggcaggagtt	cctttgggtga	cttcacagtg	aagtcttgcc	ctctctgtgg	gatatcacat	3000
cagtgggttcc	aagatttcaa	agtggtgaaa	tgaaaatgga	gcagctagta	tgttttatta	3060
ttttatgggt	cttgagtgca	tttgaagggt				3090

<210> 9
 <211> 3982
 <212> DNA
 <213> Homo Sapiens

<400> 9						
ctgggtcctg	tgtgtgccac	aggggtgggg	tgtccagcga	gcgggtctcct	cctcctgcta	60
gtgctgctgc	ggcgtccccg	ggcctccccg	agtcggggcg	gaggggagag	cgggtgtgga	120
tttgtcttga	cggtaattgt	tgcgtttcca	cgtctcggag	gcctgcgcgc	tgggttgctc	180
cttcttcggg	agcgagctgt	tctcagcgat	cccactccca	gccggggctc	cccacacaca	240
ctgggctgcg	tgcgtgtgga	gtgggacccg	cgcacacgcg	tgtctctgga	cagctacggc	300
gccgaaagaa	ctaaaattcc	agatggcaaa	ctcaatgaat	ggcagaaacc	ctgggtggtcg	360
aggaggaaat	ccccgaaaag	gtcgaatttt	gggtattatt	gatgctattc	aggatgcagt	420
tggacccctt	aagcaagctg	ccgcagatcg	caggaccgtg	gagaagactt	ggaagctcat	480
ggacaaaagt	gtaagactgt	gccaaaatcc	caaaacttcag	ttgaaaaata	gcccaccata	540
tatacttgat	attttgcctg	atacatatca	gcattttacga	cttatattga	gtaaatatga	600
tgacaaccag	aaacttgccc	aactcagtga	gaatgagtag	tttaaaatct	acattgatag	660
ccttatgaaa	aagtcaaaac	gggcaataag	actcttttaa	gaaggcaagg	agagaatgta	720
tgaagaacag	tcacaggaca	gacgaaatct	cacaaaactg	tcccttatct	tcagtcacat	780
gctggcagaa	atcaaagcaa	tctttcccaa	tgggtcaattc	cagggagata	actttcgtat	840
cacaaaagca	gatgctgctg	aattctggag	aaagtttttt	ggagacaaaa	ctatcgtacc	900
atggaaagta	ttcagacagt	gccttcatga	ggtccaccag	attagctcta	gcctggaagc	960
aatggctcta	aaatcaacaa	ttgatttaac	ttgcaatgat	tacatttcag	tttttgaatt	1020
tgatattttt	accaggctgt	ttcagccttg	gggtctctatt	ttgcggaatt	ggaatttctt	1080
agctgtgaca	catccagggt	acatggcatt	tctcacatat	gatgaagtta	aagcacgact	1140
acagaaatat	agcaccaaac	ccggaagcta	tattttccgg	ttaagttgca	ctcgattggg	1200
acagtgggcc	attggctatg	tgactgggga	tgggaatatc	ttacagacca	tacctcataa	1260
caagccctta	tttcaagccc	tgattgatgg	cagcagggaa	ggattttatc	tttatcctga	1320
tgggaggagt	tataatcctg	atttaactgg	attatgtgaa	cctacacctc	atgaccatat	1380
aaaagttaca	caggaacaat	atgaattata	ttgtgaaatg	gggtccactt	ttcagctctg	1440
taagatttgt	gcagagaatg	acaaagatgt	caagattgag	ccttgtgggc	atgtgatgtg	1500
cacctcttgc	cttacggcat	ggcaggagtc	ggatggtcag	gggtgccctt	tctgtcgttg	1560
tgaaataaaa	ggaactgagc	ccataatcgt	ggaccctttt	gatccaagag	atgaaggctc	1620
caggtgttgc	agcatcattg	accccttttg	catgccgatg	ctagacttgg	acgacgatga	1680
tgatcgtgag	gagtccttga	tgatgaatcg	gttggcaaac	gtccgaaagt	gcactgacag	1740
gcagaactca	ccagtcacat	caccaggatc	ctctcccctt	gccagagaaa	gaaagccaca	1800
gcctgaccca	ctccagatcc	cacatctaag	cctgccaccc	gtgcctcctc	gcctggatct	1860
aattcagaaa	ggcatagtta	gatctccctg	tggcagccca	acaggttcac	caaagtcttc	1920
tccttgcatg	gtgagaaaac	aagataaac	actcccagca	ccacctcctc	ccttaagaga	1980
tcctcctcca	ccgccacctg	aaagacctcc	accaatccca	ccagacaata	gactgagtag	2040
acacatccat	catgtggaaa	gcgtgccttc	cagagaccgg	ccaatgcctc	ttgaagcatg	2100
gtgccctcgg	gatgtgtttg	ggactaatca	gcttgtggga	tgtcgactcc	taggggaggg	2160
ctctccaaaa	cctggaatca	cagcgagttc	aaatgtcaat	ggaaggcaca	gtagagtggt	2220
ctctgaccca	tgctttatgc	ggaaacacag	acgccatgat	ttgccttttag	aaggagctaa	2280
ggctcttttcc	aatggtcacc	ttggaagtga	agaatatgat	gttcctcccc	ggctttctcc	2340
tcctcctcca	gttaccaccc	tcctccctag	cataaagtgt	actgggtccgt	tagcaaattc	2400
tctttcagag	aaaacaagag	accagtaga	ggaagatgat	gatgaataca	agattccttc	2460
atcccaccct	gtttccctga	attcacaacc	atctcattgt	cataatgtaa	aacctcctgt	2520
tcggctcctgt	gataatggtc	actgtatgct	gaatggaaca	catggtccat	cttcagagaa	2580


```

gaaatcaaac atccctgact taagcatata tttaaagggga gatgtttttg attcagcctc 2640
tgatcccgtg ccattaccac ctgccaggcc tccaactcgg gacaatccaa agcatgggtc 2700
ttcactcaac aggacgccct ctgattatga tcttctcatc cctccattag gtgaagatgc 2760
ttttgatgcc ctccctccat ctctcccacc tccccacact cctgcaaggc atagtctcat 2820
tgaacattca aaacctcctg gctccagtag ccggccatcc tcaggacagg atctttttct 2880
tcttcttca gatccctttg ttgatctagc aagtggccaa gttcctttgc ctctgctag 2940
aaggttacca ggtgaaaatg tcaaaactaa cagaacatca caggactatg atcagcttcc 3000
ttcatgttca gatgggttcac aggaccagc cagacccccct aaaccacgac cgcgcaggac 3060
tgcaccagaa attcaccaca gaaaaccca tgggcctgag gcggcattgg aaaatgtcga 3120
tgcaaaaatt gcaaaactca tgggagaggg ttatgccttt gaagaggtga agagagcctt 3180
agagatagcc cagaataatg tcgaagttgc ccggagcatc ctccgagaat ttgccttccc 3240
tcctccagta tccccacgtc taaatctata gcagccagaa ctgtagacac caaatggaa 3300
agcaatcgat gtattccaag agtgtggaaa taaagagaac tgagatggaa ttcaagagag 3360
aagtgtctcc tcctcgtgta gcagcttgag aagaggcttg ggagtgcagc ttctcaaagg 3420
agaccgatgc ttgctcagga tgtcgacagc tgtggcttcc ttgtttttgc tagccatatt 3480
tttaaatacag ggttgaactg acaaaaataa tttaaagacg tttacttccc ttgaactttg 3540
aacctgtgaa atgctttacc ttgtttacaa tttggcaaag ttgcagtttg ttctgtttt 3600
tagtttagtt ttgttttggt gttttgatac ctgtactgtg ttcttcacag accctttgta 3660
gcgtgggtcag gtctgctgta acatttccca ccaactctct tgctgtccac atcaacagct 3720
aaatcattta ttcatatgga tctctaccat ccccatgcct tgcccaggtc cagttccatt 3780
tctctcattc acaagatgct ttgaaggttc tgattttcaa ctgatcaaac taatgcaaaa 3840
aaaaaaaagta tgtattcttc actactgagt ttcttctttg gaaaccatca ctattgagag 3900
atgggaaaaa cctgaatgta taaagcattt atttgtcaat aaactgcctt ttgtaagggg 3960
ttttcacata aaaaaaaaaa aa 3982

```

```

<210> 10
<211> 17
<212> PRT
<213> Homo Sapiens

```

```

<400> 10
Arg Lys Pro Arg Asn Ala Gln Pro Lys Val Arg Leu Leu Lys Gly Asn
  1                      5                      10                      15
Ser

```